

10/663, 173

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal204bxd

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 JUL 12 BEILSTEIN enhanced with new display and select options,
resulting in a closer connection to BABS
NEWS 4 AUG 02 IFIPAT/IFIUDB/IFICDB reloaded with new search and display
fields
NEWS 5 AUG 02 CAPLUS and CA patent records enhanced with European and Japan
Patent Office Classifications
NEWS 6 AUG 02 The Analysis Edition of STN Express with Discover!
(Version 7.01 for Windows) now available
NEWS 7 AUG 27 BIOCOMMERCE: Changes and enhancements to content coverage
NEWS 8 AUG 27 BIOTECHABS/BIOTECHDS: Two new display fields added for legal
status data from INPADOC
NEWS 9 SEP 01 INPADOC: New family current-awareness alert (SDI) available
NEWS 10 SEP 01 New pricing for the Save Answers for SciFinder Wizard within
STN Express with Discover!
NEWS 11 SEP 01 New display format, HITSTR, available in WPIDS/WPINDEX/WPIX
NEWS 12 SEP 27 STANDARDS will no longer be available on STN
NEWS 13 SEP 27 SWETSCAN will no longer be available on STN
NEWS 14 OCT 28 KOREAPAT now available on STN

NEWS EXPRESS OCTOBER 29 CURRENT WINDOWS VERSION IS V7.01A, CURRENT
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 11 AUGUST 2004
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 21:44:25 ON 14 NOV 2004

=> fil reg

COST IN U.S. DOLLARS

SINCE FILE
ENTRY

TOTAL
SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'REGISTRY' ENTERED AT 21:44:32 ON 14 NOV 2004
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2004 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file
provided by InfoChem.

STRUCTURE FILE UPDATES: 12 NOV 2004 HIGHEST RN 780001-49-2
DICTIONARY FILE UPDATES: 12 NOV 2004 HIGHEST RN 780001-49-2

TSCA INFORMATION NOW CURRENT THROUGH MAY 21, 2004



Please note that search-term pricing does apply when
conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more
information enter HELP PROP at an arrow prompt in the file or refer
to the file summary sheet on the web at:
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>

Uploading C:\Program Files\Stnexp\Queries\10663173.str



chain nodes :

17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35

ring nodes :

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16

chain bonds :

1-25 2-24 3-23 4-22 7-18 8-28 9-27 10-16 11-29 12-17 13-21 13-32 14-19
14-20 15-30 15-31 16-26 32-33 32-34 34-35

ring bonds :

1-2 1-6 2-3 3-4 4-5 5-6 5-13 6-16 7-8 7-12 8-9 9-10 10-11 11-12 13-14
14-15 15-16

exact/norm bonds :

5-13 6-16 13-14 13-32 14-15 15-16 32-34 34-35

exact bonds :


1-25 2-24 3-23 4-22 7-18 8-28 9-27 10-16 11-29 12-17 13-21 14-19 14-20
 15-30 15-31 16-26 32-33
 normalized bonds :
 1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-12 8-9 9-10 10-11 11-12

Match level :

1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
 11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:CLASS 18:CLASS 19:CLASS
 20:CLASS 21:CLASS 22:CLASS 23:CLASS 24:CLASS 25:CLASS 26:CLASS 27:CLASS
 28:CLASS 29:CLASS 30:CLASS 31:CLASS 32:CLASS 33:CLASS 34:CLASS 35:CLASS

L1 STRUCTURE UPLOADED

=> d query



Structure attributes must be viewed using STN Express query preparation.

=> s l1

SAMPLE SEARCH INITIATED 21:44:49 FILE 'REGISTRY'
 SAMPLE SCREEN SEARCH COMPLETED - 1918 TO ITERATE

52.1% PROCESSED 1000 ITERATIONS
 INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
 SEARCH TIME: 00.00.01

0 ANSWERS

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
 BATCH **COMPLETE**
 PROJECTED ITERATIONS: 35733 TO 40987
 PROJECTED ANSWERS: 0 TO 0

L2 0 SEA SSS SAM L1

=> s l1 full

FULL SEARCH INITIATED 21:44:52 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 38243 TO ITERATE

100.0% PROCESSED 38243 ITERATIONS 12 ANSWERS
SEARCH TIME: 00.00.01

L3 12 SEA SSS FUL L1

=> fil caplus

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	155.42	155.63

FILE 'CAPLUS' ENTERED AT 21:44:55 ON 14 NOV 2004
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 14 Nov 2004 VOL 141 ISS 21
FILE LAST UPDATED: 12 Nov 2004 (20041112/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l3

L4 79 L3


=> d l4 1'-79 abs ibib hitstr

L4 ANSWER 1 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB The serotonin transporter (SERT) belongs to a family of sodium chloride-dependent transporters responsible for uptake of amino acids and biogenic amines from extracellular spaces. SERT represents the main pharmacol. target in the treatment of several clin. conditions, including depression and anxiety. Serotonin-selective reuptake inhibitors and tricyclic antidepressants are the most predominantly prescribed drugs in the treatment of depression. In addition to antidepressants also psychostimulants, like cocaine and amphetamines, are important SERT antagonists. In the present study, the authors report the cloning and characterization of chicken SERT. Although the uptake kinetic was very similar to human SERT, the pharmacol. profiles differed considerably for the two species. The authors find that chicken SERT is capable of discriminating between different serotonin-selective reuptake inhibitors; thus, the potency of S-citalopram and paroxetine is reduced more than 40-fold. A cross-species chimera strategy was undertaken and followed by species-scanning mutagenesis. Differences in pharmacol. profiles were tracked to amino acid residues 169, 172, and 586 in human SERT. Structure-activity studies on structurally related compds. indicated that species divergences in drug sensitivity between human and chicken SERT were arising from differences in coordination or recognition of an important aminomethyl pharmacophoric substructure, which is shared by all high affinity antidepressants. Consequently, the authors suggest that Ala169 and Ile172 of human SERT are important residues in sensing the N-methylation state of SERT antagonists.

ACCESSION NUMBER: 2004:791467 CAPLUS
 TITLE: The Chicken Serotonin Transporter Discriminates between Serotonin-selective Reuptake Inhibitors: A Species-scanning mutagenesis study
 AUTHOR(S): Larsen, Mads Areum; Elfvig, Betina; Wiborg, Ove
 CORPORATE SOURCE: Laboratory of Molecular Neurobiology, Department of Biological Psychiatry, Aarhus Psychiatric University Hospital, Risskov, 8240, Den.
 SOURCE: Journal of Biological Chemistry (2004), 279(40), 42147-42156
 CODEN: JBCHA3; ISSN: 0021-9258
 PUBLISHER: American Society for Biochemistry and Molecular Biology
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT INDEXING IN PROGRESS
 IT 07857-41-8, Desmethylertraline
 RL: PAC (Pharmacological activity); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (chicken serotonin transporter discriminates between serotonin-selective reuptake inhibitors, a Species-scanning mutagenesis study)
 RN 07857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 1 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L4 ANSWER 2 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB Sertraline and paroxetine are frequently prescribed SSRIs for long-term treatment of major depression. Nevertheless, continuous follow-ups of drug concns. prevailing in patients during the whole treatment period are not available. Hence, in a large phase IV clin. trial, a total of 353 patients with major depression were enrolled for a 6-mo comparison of sertraline (50-150 mg daily) and paroxetine (20-60 mg daily). The present study reports the pharmacokinetic results of up to eight serum samples per patient. 1. A profound variability was found in the interindividual steady state and trough serum levels of sertraline, desmethylertraline and paroxetine: the coefficient of variation (CV) was 59% for sertraline, 51% for desmethylertraline, 27% for the ratio desmethylertraline/sertraline (50 mg/day), and 71% for paroxetine (20 mg/day). The intraindividual CV for the ratio desmethylertraline/sertraline was only 19%, indicating intraindividual metabolizing stability over time. Both sertraline and paroxetine displayed sex differences in the dose-concentration correlation. 2. It was possible to predict sertraline, but not paroxetine, steady state levels. 3. The terminal elimination t1/2 for both drugs after 6 mo of treatments was similar to data previously reported from short-term withdrawal studies. 4. No correlation between serum drug concns. and clin. effect was detected for either sertraline or paroxetine. For the future, continuous efforts are warranted to perform PK investigations in the natural clin. setting in which the drugs are usually prescribed.

ACCESSION NUMBER: 2004:682102 CAPLUS
 DOCUMENT NUMBER: 141:342815
 TITLE: Serum disposition of sertraline, N-desmethylertraline and paroxetine: A pharmacokinetic evaluation of repeated drug concentration measurements during 6 months of treatment for major depression
 AUTHOR(S): Reis, Margareta; Aberg-Wistedt, Anna; Agren, Hans; Hoeglund, Peter; Akerblad, Ann-Charlotte; Bengtsson, Finn
 CORPORATE SOURCE: Department of Medicine and Care, Division of Clinical Pharmacology, Faculty of Health Sciences, Linköping University, Linköping, Swed.
 SOURCE: Human Psychopharmacology (2004), 19(5), 283-291
 CODEN: HUPSEC; ISSN: 0885-6222
 PUBLISHER: John Wiley & Sons Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 07857-41-8, N-Desmethylertraline
 RL: PKT (Pharmacokinetics); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (serum disposition of sertraline, desmethylertraline and paroxetine)
 RN 07857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 2 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L4 ANSWER 3 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN

AB The 1st indication that select antidepressants and their metabolites accumulate in organisms residing in effluent-dominated streams was provided. Fluoxetine, norfluoxetine, sertraline, and desmethylsertraline were detected in brain, liver, and lateral muscle tissues of 3 fish populations, *Lepomis macrochirus* (bluegill), *Ictalurus punctatus* (channel catfish) and *Pomoxis nigromaculatus* (black crappie). The highest concns. of all compds. were observed in brain and liver; lowest concns. were detected

in lateral muscle. Fish with larger body size such as *I. punctatus* and

P. nigromaculatus generally accumulated higher amts. of all the compds. tested.

ACCESSION NUMBER: 2004:613535 CAPLUS.

DOCUMENT NUMBER: 141:230101

TITLE: Determination of select antidepressants in organisms from a municipal effluent-dominated stream

AUTHOR(S): Brooks, Bryan W.; Chambliss, C. Kevin; Stanley, Jacob K.; Ramirez, Alejandro; Banks, Kenneth; Johnson, Robert D.; Lewis, Russell J.

CORPORATE SOURCE: Department of Environmental Studies, Center for Reservoir and Aquatic Systems Research, Baylor University, Waco, TX, 76798, USA

SOURCE: Preprints of Extended Abstracts presented at the ACS National Meeting, American Chemical Society, Division of Environmental Chemistry (2004), 44(2), 1351-1355

PUBLISHER: American Chemical Society, Division of Environmental Chemistry

DOCUMENT TYPE: Journal; (computer optical disk)

LANGUAGE: English


IT 87857-41-8, Desmethylsertraline

RL: POL (Pollutant); OCCU (Occurrence) (antidepressants in fish from municipal effluent-dominated streams)

RN 87857-41-8 CAPLUS

CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE

FORMAT

L4 ANSWER 3 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

L4 ANSWER 4 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN

AB The invention provides compns. and methods for the treatment of a vaso-occlusive event. More particularly, the invention provides a combination therapy for the treatment of a vaso-occlusive event

comprising the administration to a subject of a selective serotonin reuptake inhibitor in combination with a cyclooxygenase-2 selective inhibitor.

ACCESSION NUMBER: 2004:565129 CAPLUS

DOCUMENT NUMBER: 141:99705

TITLE: Methods and compositions using cyclooxygenase-2 selective inhibitors and selective serotonin reuptake inhibitors for the treatment or prevention of a vaso-occlusive event

INVENTOR(S): Stephenson, Diane T.; Taylor, Duncan P.

PATENT ASSIGNEE(S): Pharmacia Corporation, USA

SOURCE: PCT Int. Appl., 159 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004058354	A1	20040715	WO 2003-US40955	20031222
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HK, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ				
RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, NI, SN, TD, TG				
US 2004171664	A1	20040902	US 2003-743485	20031222
PRIORITY APPLN. INFO.:			US 2002-435078P	P 20021220

OTHER SOURCE(S): MARPAT 141:99705

IT 87857-41-8, N-Desmethylsertraline

RL: POC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)

(cyclooxygenase 2 selective inhibitors and selective serotonin reuptake


inhibitors for treatment or prevention of vaso-occlusive event)

RN 87857-41-8 CAPLUS

CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 4 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



L4 ANSWER 5 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
AB Background: Selective serotonin reuptake inhibitor antidepressants (SSRIs) and benzodiazepines are frequently used to treat maternal depression and anxiety disorders during pregnancy. Recent reports suggest that prenatal SSRI exposure is associated with a neonatal discontinuation syndrome. It remains unclear whether these symptoms are directly related to SSRI exposure alone or are due to concurrent pharmacol. factors. Also, this study explores relationships between neonatal outcomes and medication levels during pregnancy, at delivery, and in the newborn period. Method: This study sought to compare newborn behavior following second and third trimester exposure to either single-agent SSRIs (group 1) or SSRIs combined with clonazepam (group 2). A prospective cohort of mothers and their infants (N = 46) who had received SSRI medication alone or in combination with clonazepam were studied from June 1996 through June 2000 and compared with a non-exposed control group (N = 23). Infants were assessed in the newborn period for signs suggestive of a "discontinuation syndrome". Maternal drug levels were measured during the pregnancy and at delivery. Infant drug levels from cord blood and at day 2 of life

were also obtained. Results: Overall, 30% of the exposed infants (groups 1 and 2, N = 14) showed symptoms of transient poor neonatal adaptation compared with 9% (N = 2) of control infants. In group 1, 25% had symptoms (fluoxetine N = 3; paroxetine N = 3; sertraline N = 1) and in group 2, 39% of infants had symptoms (paroxetine with clonazepam, N = 7). Symptoms were typically mild respiratory distress and, less commonly, hypotonia. Symptoms were self limited and not associated with other neonatal conditions. When paroxetine was combined with clonazepam, infants with symptoms had significantly elevated paroxetine levels when compared with similarly exposed infants without symptoms (p < .05). Among single-agent paroxetine-exposed infants, drug levels did not differ significantly between those with and without symptoms. Maternal dose of clonazepam was significantly higher (p < .05) during pregnancy and at delivery among symptomatic infants compared with non-symptomatic infants. Developmental outcomes at 2 and 8 mo of age did not differ between symptomatic and non-symptomatic infants. Conclusion: While transient neonatal symptoms were found in infants after single-agent prenatal exposure to SSRIs and when paroxetine was combined with clonazepam, the addition of clonazepam appeared to alter paroxetine metabolism, leading to increased drug levels and risk for transient neonatal symptoms. These data highlight the importance of accounting for a variety of pharmacol. factors beyond single-agent SSRI exposure that may lead to poor neonatal adaptation. Further studies are needed with a larger sample of infants to determine the role of clonazepam and whether similar outcomes occur when exposure includes other SSRIs in combination with clonazepam.


ACCESSION NUMBER: 2004:281036 CAPLUS
DOCUMENT NUMBER: 140:314965
TITLE: Pharmacologic factors associated with transient neonatal symptoms following prenatal psychotropic medication exposure
AUTHOR(S): Oberlander, Tim F.; Misri, Shaila; Fitzgerald, Colleen
E.; Kostaras, Xanthoula; Rurak, Dan; Riggs, Wayne

L4 ANSWER 6 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
AB Trans-, (1R,4S)-, and (1S,4R)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenamines are claimed. In functional monoamine uptake inhibition assays for serotonin, norepinephrine, and dopamine, the (1R,4S)- isomer showed IC50 = 0.0077, 0.0096, and 0.0064 µM, resp.
ACCESSION NUMBER: 2004:252469 CAPLUS
DOCUMENT NUMBER: 140:287185
TITLE: Preparation of trans-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenamine for treatment of CNS disorders.
INVENTOR(S): Jerussi, Thomas P.; Fang, Qun Kevin; Currie, Mark
PATENT ASSIGNEE(S): Sepracor, Inc., USA
SOURCE: PCT Int. Appl., 36 pp.
DOCUMENT TYPE: CODEN: PIXXD2
LANGUAGE: Patent
FAMILY ACC. NUM. COUNT: English
PATENT INFORMATION: 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004024669	A1	20040325	WO 2003-US29110	20030916
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZH, ZW, AM, AZ, BY, KG, KZ, MD				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, NG, SN, TD, TG				
US 2004092605	A1	20040513	US 2003-663173	20030916
PRIORITY APPLN. INFO.:			US 2002-411304P	P 20020916
			US 2002-411305P	P 20020916


OTHER SOURCE(S): CASREACT 140:287185
IT 675126-04-2P, trans-4-(3,4-Dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenamine 675126-05-3P, (1R,4S)-4-(3,4-Dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenamine 675126-06-4P, (1S,4R)-4-(3,4-Dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenamine 675126-07-5P, (1S,4R)-4-(3,4-Dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenamine hydrochloride 675126-08-6P, (1R,4S)-4-(3,4-Dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenamine hydrochloride 675126-09-7P, (1R,4R)-4-(3,4-Dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenamine hydrochloride 675126-10-0P, (1S,4S)-4-(3,4-Dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenamine hydrochloride
RI: PAC (Pharmacological activity); BIOL (Biological study); PREP (Preparation); USES (Uses)
(preparation of dichlorophenyltetrahydronaphthalenamines for treatment of CNS disorders)
RN 675126-04-2 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1R,4S)-rel- (9CI) (CA INDEX NAME)

L4 ANSWER 5 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
CORPORATE SOURCE: Department of Pediatrics, University of British Columbia, Vancouver, BC, Can.
SOURCE: Journal of Clinical Psychiatry (2004), 65(2), 230-237
CODEN: JCLPDE; ISSN: 0160-6689
PUBLISHER: Physicians Postgraduate Press, Inc.
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 67857-41-8, Desmethylertraline
RL: BSU (Biological study, unclassified); BIOL (Biological study) (pharmacol. factors associated with transient neonatal symptoms following prenatal psychotropic medication exposure)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)- (9CI) (CA INDEX NAME)
Absolute stereochemistry.




REFERENCE COUNT: 36
THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE
FORMAT

L4 ANSWER 6 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
Relative stereochemistry.
RN 675126-05-3 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1R,4S)- (9CI) (CA INDEX NAME)
Absolute stereochemistry.




RN 675126-06-4 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4R)- (9CI) (CA INDEX NAME)
Absolute stereochemistry.



RN 675126-07-5 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, hydrochloride, (1S,4R)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



● HCl

RN 675126-08-6 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, hydrochloride, (1R,4S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



● HCl

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE


FORMAT



● HCl

RN 675126-09-7 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, hydrochloride, (1R,4R)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



● HCl

RN 675126-10-0 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, hydrochloride, (1S,4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 7 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
AB Methods and compns. for reducing or controlling appetite using cyclooxygenase-2 inhibitors are disclosed.
ACCESSION NUMBER: 2004:100963 CAPLUS
DOCUMENT NUMBER: 140:139550
TITLE: Cyclooxygenase-2 inhibitors for appetite suppression
INVENTOR(S): Schwartz, Yael
PATENT ASSIGNEE(S): Sepracor Inc., USA
SOURCE: PCT Int. Appl., 10 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004010955	A2	20040205	WO 2003-US23922	20030731
WO 2004010955	A3	20040805		


W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, D2, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.: US 2002-39976P P 20020731


IT 87857-41-8
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(cyclooxygenase-2 inhibitors for appetite suppression, and use with other agents)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 8 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB The high incidence of psychiatric illness in the postpartum period and the increasing percentage of women who breastfeed has focused attention on the treatment of breastfeeding women with psychotropic medications and, addnl., the exposure of nursing infants to these medications. Consequently, there has been an increased effort to develop standardized methods for quantifying psychotropic medications in breast milk. This paper details a novel method for quantifying the concns. of multiple selective serotonin reuptake inhibitors (SSRIs) and tricyclic antidepressants (TCAs) in breast milk. The method consists of a common liquid/liquid and solid-phase extraction followed by HPLC separation on a common column and UV detection. Assay system 1 measures fluoxetine, norfluoxetine, fluvoxamine, and paroxetine; assay 2 measures sertraline and desmethylsertraline; and assay 3 measures the TCAs including doxepin, nortriptyline, desipramine, imipramine, nortriptyline, and amitriptyline. The method is shown to be a highly accurate and precise technique for measuring 12 different antidepressants in human breast milk and to be free of the matrix effects often encountered in breast milk drug analyses.

ACCESSION NUMBER: 2004:42859 CAPLUS
 DOCUMENT NUMBER: 141:167098
 TITLE: A Novel System for the Determination of Antidepressant Concentrations in Human Breast Milk
 AUTHOR(S): Hostetter, Amy L.; Stowe, Zachary N.; Cox, Mary; Ritchie, James C.
 CORPORATE SOURCE: Department of Psychiatry and Behavioral Sciences, Emory University School of Medicine, Atlanta, GA, 30322, USA
 SOURCE: Therapeutic Drug Monitoring (2004), 26(1), 47-52
 CODEN: TDMODV; ISSN: 0163-4356
 PUBLISHER: Lippincott Williams & Wilkins
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylsertraline
 RL: ANT (Analyte); THU (Therapeutic use); ANST (Analytical study); BIOL (Biological study); USES (Uses)
 (assay system II of novel technique consisting liquid/liquid and solid-phase extraction followed by HPLC separation and UV detection accurately and precisely determined concentration of desmethylsertraline in human breast milk)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)
 Absolute stereochemistry.




REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L4 ANSWER 9 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB Introduction: Selective serotonin reuptake inhibitors (SSRIs) are popularly prescribed for treating depression. With a few exceptions, these psychotropic medications are not approved by aeromedical regulatory authorities for use by aviators. Since SSRIs have the potential for impairing performance and causing drug-drug interactions, the prevalence of SSRIs in pilot fatalities of civil aviation accidents was evaluated. Methods: Postmortem samples from pilots involved in fatal civil aircraft accidents are submitted to the Civil Aerospace Medical Institute (CAMI) for toxicol. evaluation. Findings from such evaluations are maintained in the CAMI Toxicol. Database. This database was examined for the presence of SSRIs in pilot fatalities of the accidents that occurred during 1990-2001. Results: Out of 4,184 fatal civil aviation accidents from which CAMI received samples, there were 61 accidents in which pilot fatalities had SSRIs. Of these accidents, 56 were of the general aviation category, 2 were of the air taxi and commuter category, 2 were of the agricultural category, and 1 was of the ultralight category. Blood concns. of SSRIs in the fatalities were 11-1121 ng · ml⁻¹ for fluoxetine; 47-13102 ng · ml⁻¹ for sertraline; 68-1441 ng · ml⁻¹ for paroxetine; and 314-462 ng · ml⁻¹ for citalopram. In 39 of the 61 pilots, other drugs-for example, analgesics, anti histaminics, benzodiazepines, narcotic analgesics, and/or sympathomimetics-and/or ethanol were also present. As determined by the National Transportation Safety Board, the use of an SSRI [With or without other drugs) and/or ethanol] has been a contributory factor in at least 9 of the 61 accidents. Conclusions: Nos. of SSRI-involved accidents were low, and blood SSRI concns. in the associated pilot fatalities ranged from subtherapeutic to toxic levels. However, the interactive effects of other drug(s), ethanol, and/or even altitude hypoxia in producing adverse effects in the pilots cannot be ruled out. Findings from this study should be useful in investigating SSRI and other substance-involved accidents and in making decisions concerning the use of SSRIs in aviation.

ACCESSION NUMBER: 2003:988335 CAPLUS
 DOCUMENT NUMBER: 141:2479
 TITLE: Selective serotonin reuptake inhibitors in pilot fatalities of civil aviation accidents, 1990-2001
 AUTHOR(S): Akin, Ahmet; Chaturvedi, Arvind K.
 CORPORATE SOURCE: Bioaeronautical Sciences Research Laboratory, Aerospace Medical Research Division, Civil Aerospace Medical Institute, U.S. Department of Transportation, Federal Aviation Administration, Oklahoma City, USA
 SOURCE: Aviation, Space and Environmental Medicine (2003), 74(11), 1169-1176
 CODEN: ASEMCG; ISSN: 0095-6562
 PUBLISHER: Aerospace Medical Association
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylsertraline
 RL: ADV (Adverse effect, including toxicity); ANT (Analyte); ANST (Analytical study); BIOL (Biological study)
 (serotonin reuptake inhibitors in pilot fatalities of civil aviation


L4 ANSWER 9 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
accidents, 1990-2001)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR
THIS
FORMAT
RECORD. ALL CITATIONS AVAILABLE IN THE RE

L4 ANSWER 10 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



L4 ANSWER 10 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
AB The invention discloses pharmacol. methods for the prevention of
amelioration of sleep-related breathing disorders via administration of
agents or combinations of agents that possess serotonin-related
pharmacol.
activity. Agents of the invention include e.g. ondansetron.
ACCESSION NUMBER: 2003:532342 CAPLUS
DOCUMENT NUMBER: 139:95476
TITLE: Agents having serotonin-related pharmacol. activity
for the pharmacological treatment of sleep apnea and
other sleep-related breathing disorders
Radulovacki, Miodrag; Carley, David W.
INVENTOR(S):
PATENT ASSIGNEE(S): USA
SOURCE: U.S. Pat. Appl. Publ., 20 pp., Cont.-in-part of U.S.
Ser. No. 16,901.
CODEN: USXXCO
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 2003130266	A1	20030710	US 2002-285277	20021031
US 2002086870	A1	20020704	US 2001-16901	20011214
US 6727242	B2	20040427		
WO 2004041272	A2	20040521	WO 2003-US34592	20031029
WO 2004041272	A3	20040916		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GR, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, ML, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GN, GW, GQ, GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO: US 2001-16901 A2 20011214
US 1998-76216P P 19980227
WO 1999-US4347 W 19990226
US 2000-622823 A1 20000823
US 2002-285277 A 20021031


IT 87857-41-8, Demethylsertraline
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(agents with serotonin-related pharmacol. activity for treatment of sleep apnea and other sleep-related breathing disorders)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 11 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
AB Pharmacokinetic drug-drug interactions often occur at the level of P-glycoprotein (Pgp). To study possible interactions caused by the newer antidepressants we investigated citalopram, fluoxetine, fluvoxamine, paroxetine, reboxetine, sertraline, and venlafaxine and their major metabolites desmethylcitalopram, norfluoxetine, paroxetine-metabolite (paroxetine-M), desmethylsertraline, N-desmethylvenlafaxine, and O-desmethylvenlafaxine for their ability to inhibit Pgp. Pgp inhibition was studied by a fluorometric assay using calcein-acetoxymethyl ester as Pgp substrate and two different cell systems: L-MDR1 cells (model for human Pgp) and primary porcine brain capillary endothelial cells (pBCECs, model for the blood-brain barrier). Both cell systems proved to be suitable for the evaluation of Pgp inhibitory potency of drugs. All antidepressants tested except O-desmethylvenlafaxine showed Pgp inhibitory activity with sertraline, desmethylsertraline, and paroxetine being the most potent, comparable with the well known Pgp inhibitor quinidine. In L-MDR1 cells fluoxetine, norfluoxetine, fluvoxamine, reboxetine, and paroxetine-M revealed intermediate Pgp inhibition and citalopram, desmethylcitalopram, venlafaxine, and N-desmethylvenlafaxine were only weak inhibitors. The ranking order was similar in pBCECs. The fact that some of the compds. tested exert Pgp inhibitor effects at similar concns. as quinidine suggests that pharmacokinetic drug-drug interactions between the newer antidepressants and Pgp substrates should now be thoroughly studied in vivo.

ACCESSION NUMBER: 2003:262485 CAPLUS
DOCUMENT NUMBER: 139:207091
TITLE: Inhibition of P-glycoprotein by newer antidepressants
AUTHOR(S): Weiss, Johanna; Dormann, Sven-Maria Gregor; Martin-Facklam, Meret; Kerpen, Christian Johannes; Ketabi-Kiyanvash, Nahal; Haefeli, Walter Emil
CORPORATE SOURCE: Department of Internal Medicine VI, Clinical Pharmacology, and Pharmacoeconomics, University of Heidelberg, Heidelberg, Germany
SOURCE: Journal of Pharmacology and Experimental Therapeutics (2003), 305(1), 197-204
CODEN: JPETAR; ISSN: 0022-3565
PUBLISHER: American Society for Pharmacology and Experimental Therapeutics
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: ADV (Adverse effect, including toxicity); PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(antidepressant inhibition of P-glycoprotein in relation to pharmacokinetic drug interactions)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT:
THIS
FORMAT

40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE

IT 87857-41-8, Desmethylertraline
RL: PKT (Pharmacokinetics); BIOL (Biological study)
(pharmacokinetics of sertraline excretion into human breast milk as
predictors of infant serum concns.)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT:
THIS
FORMAT

27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE

AB Background: The purpose of this study was to attain a new landmark in the
area of selective serotonin reuptake inhibitor therapy during lactation

by

establishing a basis for interpreting infant serum concns. and for
minimizing infant exposure in the absence of treatment-emergent side
effects. Method: Breast milk and paired maternal and infant sera were
collected following maternal treatment with sertraline monotherapy

(25-200
mg/day) administered once daily. Sertraline and its major metabolite

were

measured in breast milk and serum samples using high-performance liquid
chromatog. with UV detection (limit of detection = 2 ng/mL). Results:
Twenty-six nursing women with DSM-IV major depressive disorder
participated in the study; the mean (SD) daily sertraline dose was 123.9
(62.8) mg/day. Fifteen women submitted 182 breast milk samples for anal.
of gradient (foremilk to hindmilk) and time course of medication
excretion. The milk/plasma ratio was highly variable (range, 0.42-4.81).
A significant gradient and time course of excretion for both sertraline

(p

< .001 for both) and desmethylertraline (p < .001 for gradient and p <
.046 for time course) were observed, with the highest concns. observed

in the

hindmilk 8 to 9 h after maternal ingestion. Math. modeling of sertraline
and desmethylertraline excretion revealed that discarding breast milk 9

h

after maternal dose decreased the infant daily dose of sertraline by a
mean of 17.1% (1.8%). Twenty-two mother/infant sera pairs were obtained.
Sertraline was detectable in 4 infants (18% of sample), and
desmethylertraline was found in 11 infants (50% of sample). The mean
(SD) maximum calculated nursing infant dose of sertraline, 0.67 (0.61)

mg/day,

and desmethylertraline, 1.44 (1.36) mg/day, represented 0.54% (0.49%) of
the maternal daily dose. The maximum infant dose of desmethylertraline

(p <

.002) significantly correlated with infant serum desmethylertraline
concns. (ng/mL). In contrast, maternal daily dose, duration of

medication

exposure, and infant age and weight at sampling did not correlate with

either

detectability (< 2 ng/mL vs. ≥ 2 ng/mL) or absolute concns. (ng/mL) in
infant serum. No adverse events were reported or documented in any
infant. Conclusion: These results extend previous studies by
demonstrating the utility of breast milk anal. in interpreting infant
serum concns. and minimizing infant exposure.

ACCESSION NUMBER: 2003:161587 CAPLUS

DOCUMENT NUMBER: 138:297064

TITLE: The pharmacokinetics of sertraline excretion into
human breast milk: determinants of infant serum
concentrations

AUTHOR(S): Stowe, Zachary N.; Hostetter, Amy L.; Owens, Michael
J.; Ritchie, James C.; Sternberg, Kevan; Cohen, Lee
S.; Nemeroff, Charles B.

CORPORATE SOURCE: Department of Psychiatry and Behavioral Sciences,
Emory University School of Medicine, Atlanta, GA, USA

SOURCE: Journal of Clinical Psychiatry (2003), 64(1), 73-80
CODEN: JCLPDE; ISSN: 0160-6689

PUBLISHER: Physicians Postgraduate Press, Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The main purpose of this study was the investigation of quant.
structure-activity relationships of serotonin transporter ligands with
regard to the future development of potential new and selective PET
radiotracers for the serotonin transporter. A heterogeneous data set of
19 selective and non-selective serotonin reuptake inhibitors was used.
Affinity data for both the serotonin transporter and the norepinephrine
transporter was available. As a necessary prerequisite for our 3D QSAR
studies a reasonable alignment of the compds. was developed using GASF.
It was based on an existing pharmacophore model. In addition to the
widely

used CoMFA method, the somewhat newer CoMSIA method was applied.
Statistically reliable CoMFA models for both the serotonin transporter
(q²=0.538) and the norepinephrine transporter (q²=0.445) were developed,
further improving the internal predictability by applying region focusing
for the serotonin transporter (q²=0.674). These models were compared

with

the CoMSIA models for the serotonin and the norepinephrine transporter
that yielded comparable cross-validated correlation coeffs. (q²=0.531 and
q²=0.502, resp.). Certain structural features that are distinctive of
each transporter and important for high binding affinity were identified.
Highly comparable results were obtained for CoMFA and CoMSIA. Both
methods were applied to elucidate structural requirements for serotonin
transporter selectivity. The resulting CoMSIA map provides important
information for lead optimization with respect to selectivity

enhancement.

ACCESSION NUMBER: 2003:109510 CAPLUS

DOCUMENT NUMBER: 139:159428

TITLE: 3D QSAR of serotonin transporter ligands: CoMFA and
CoMSIA studies

AUTHOR(S): Wellisaw, Julia; Machulla, Hans-Juergen; Kovar,
Karl-Artur

CORPORATE SOURCE: Pharmaceutical Institute, University of Tübingen,
Tübingen, 72076, Germany

SOURCE: Quantitative Structure-Activity Relationships (2002),
21(6), 577-589

CODEN: QSARDI; ISSN: 0931-8771

PUBLISHER: Wiley-VCH Verlag GmbH & Co. KGaA

DOCUMENT TYPE: Journal

LANGUAGE: English

IT 87857-41-8, Desmethylertraline

RL: PAC (Pharmacological activity); BIOL (Biological study)


(QSAR of serotonin transporter ligands)

RN 87857-41-8 CAPLUS

CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 13 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT:
THIS
FORMAT

42 THERE ARE 42 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE


L4 ANSWER 14 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN

AB Separating drugs which are both polar and basic has long been difficult because of the limited operating pH range of conventional HPLC columns. This paper describes a liquid chromatog. method capable of being used with either diode-array or mass spectrometric detection for the identification and quantitation of 10 antidepressant and 2 antipsychotic drugs, all of which have serotonergic activity. In developing the method, the effects of varying buffers and mobile phase pH and of adding modifying agents on resolution and capacity factors were investigated. The organic buffers ammonia, glycine, and triethylamine were each used in a mobile phase made up of 32.5% buffer/67.5% methanol (volume/volume) at a pH of 10.0. Addnl., 4 different concns. each of THF and acetonitrile were added to investigate the effect of a modifying agent on resolution and retention. In general, decreasing mobile phase pH reduced retention times and decreased resolution. Adding THF in place of the same amount of methanol tended to decrease retention times, and adding acetonitrile tended to slightly increase retention times. However, addition of both marginally improved resolution. This method was used to satisfactorily analyze brain, blood, liver, urine, vitreous fluid, and stomach contents in subjects known to have used these drugs. (c) 2003 Preston Publications.

ACCESSION NUMBER: 2003:70640 CAPLUS
DOCUMENT NUMBER: 138:315995
TITLE: LC-MS analysis of serotonergic drugs
AUTHOR(S): Goeringer, Kabrena E.; McIntyre, Iain M.; Drummer, Olaf H.
CORPORATE SOURCE: Victorian Institute of Forensic Medicine and Department of Forensic Medicine, Monash University, Southbank, 3006, Australia
SOURCE: Journal of Analytical Toxicology (2003), 27(1), 30-35
CODEN: JATOD3; ISSN: 0146-4760
PUBLISHER: Preston Publications
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylesterline
RL: ANT (Analyte): ANST (Analytical study)
(LC-MS anal. of serotonergic drugs)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 14 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT:
THIS
FORMAT

24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE


L4 ANSWER 15 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN

AB Bile is, in certain cases, collected together with blood from different sites (heart, brain, femoral), urine and other organs or matrices. This study reports comparative results obtained from the anal. of blood and bile for different drugs found: acetaminophen, amphetamine and related compds., several antidepressants, several benzodiazepines, cocaine and its metabolites, dextropropoxyphene and its metabolite, hydroxyzine, methadone and metabolite, morphine and codeine, levomepromazine, thioridazine, propranolol, tramadol and its metabolite. Several findings are presented: There were no significant differences in the levels of the compds. among the samples of blood obtained from different sites. Levels in bile are generally several fold higher than those in blood. The mean bile to blood ratios vary from about 1 (for acetaminophen, amphetamine) to about 2000 (for desmethylobazam). In certain cases (16 over 44), although the drug or its metabolite was not detected in blood from different sites, it was detected in bile. As other authors had advocated, it is very useful to ask the pathologist to take the gall bladder with its contents together, with the other samples, in order that the sample of bile can be used in the comprehensive toxicol. anal. and therefore be complementary to the other fluids or matrices. An addnl. advantage for using bile is that the concns. of drugs or their metabolites are generally several fold higher than their blood concns.

ACCESSION NUMBER: 2002:669216 CAPLUS
DOCUMENT NUMBER: 138:164923
TITLE: Bile analysis of drugs in postmortem cases
AUTHOR(S): Vanbinst, R.; Koenig, J.; Di Fazio, V.; Hassoun, A.
CORPORATE SOURCE: St-Luc Hospital, Laboratory of Toxicology, Universite Catholique de Louvain, Brussels, 1200, Belg.
SOURCE: Forensic Science International (2002), 128(1-2), 35-40
CODEN: FSINDR; ISSN: 0379-0738
PUBLISHER: Elsevier Science Ireland Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylesterline
RL: BSU (Biological study, unclassified); BIOL (Biological study)
(drug screening in bile compared with blood, urine, and gastric content in forensic postmortem cases)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 15 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT:
THIS
FORMAT

16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE

L4 ANSWER 16 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN


AB Most drugs are excreted into breast milk to some extent and are bioavailable to the infant. The ability to predict the approx. amount of drug that might be present in milk from the drug structure would be very useful in the clin. setting. The aim of this research was to simplify and upgrade the previously developed model for prediction of the milk to plasma (M/P) concentration ratio, given only the mol. structure of the drug. The set of 123 drug compds., with exptl. derived M/P values taken from the literature, was used to develop, test and validate a predictive model. Each compound was encoded with 71 calculated mol. structure descriptors, including constitutional descriptors, topol. descriptors, mol. connectivity, geometrical descriptors, quantum chemical descriptors, physicochem. descriptors and liquid properties. Genetic algorithm was used to select a subset of the descriptors that best describe the drug transfer into breast milk and artificial neural network (ANN) to correlate selected descriptors with the M/P ratio and develop a QSAR. The averaged literature M/P values were used as the ANN's output and calculated mol. descriptors as the inputs. A nine-descriptor nonlinear computational neural network model has been developed for the estimation of M/P ratio values for a data set of 123 drugs. The model included the percent of oxygen, parachor, d., HOMO energy (HOMO), topol. indexes (χV_2 , χ_2 and χ_1) and shape indexes (κ_3 , κ_2), as the inputs had four hidden neurons and one output neuron. The QSPR that was developed indicates that mol. size (parachor, d.) shape (topol. shape indexes, mol. connectivity indexes) and electronic properties (HOMO) are the most important for drug transfer into breast milk. Unlike previously reported models, the QSPR model described here does not require exptl. derived parameters and could potentially provide a useful prediction of M/P ratio of new drugs only from a sketch of their structure and this approach might also be useful for drug information service. Regardless of the model or method used to estimate drug transfer into breast milk, these predictions should only be used to assist in the evaluation of risk, in conjunction with assessment of the infant's response.

ACCESSION NUMBER: 2002:438488 CAPLUS
DOCUMENT NUMBER: 138:32765
TITLE: Molecular descriptors that influence the amount of drugs transfer into human breast milk
AUTHOR(S): Agatonovic-Kustrin, S.; Ling, L. H.; Tham, S. Y.; Alany, R. G.
CORPORATE SOURCE: School of Pharmaceutical, Molecular and Biomedical Science, University of South Australia, Adelaide, 5000, Australia
SOURCE: Journal of Pharmaceutical and Biomedical Analysis (2002), 29(1-2), 103-119
CODEN: JPBADA; ISSN: 0731-7085
PUBLISHER: Elsevier Science B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8
RL: PKT (Pharmacokinetics); PRP (Properties); BIOL (Biological study) (mol. descriptors that influence the amount of drugs transfer into human breast milk in relation to their bioavailability to the newborn)

L4 ANSWER 16 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)

RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT:
THIS
FORMAT


163 THERE ARE 163 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE

L4 ANSWER 17 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN

AB LC/MS/MS provides superior sensitivity and selectivity, rapid anal., maximized development efficiencies, and improved turnaround times. The challenges of large-scale LC/MS/MS anal. may be overcome with careful planning and the suggested troubleshooting techniques.

ACCESSION NUMBER: 2002:284366 CAPLUS
DOCUMENT NUMBER: 137:121824
TITLE: Systematic troubleshooting for LC/MS/MS
AUTHOR(S): Maidong, Weng; Halls, Timothy D. J.
CORPORATE SOURCE: Covance Laboratories Inc., Madison, WI, 53704, USA
SOURCE: Pharmaceutical Technology North America (2002), 26(3), 102,104,106,108,110,112,114,116,118,120
CODEN: PTNABQ; ISSN: 1534-2131
PUBLISHER: Advanstar Communications, Inc.
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylertraline
RL: ANT (Analyte); ANST (Analytical study) (anal. of drugs in blood and urine by LC/MS/MS and troubleshooting techniques)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT:
THIS
FORMAT

24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE

L4 ANSWER 18 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
AB LC/MS/MS provides superior sensitivity and selectivity, rapid anal., maximized development efficiencies, and improved turnaround times. The challenges of large-scale LC/MS/MS anal. may be overcome with careful planning and the suggested troubleshooting techniques. Part 1 presents troubleshooting techniques related to sample preparation and chromatog.


ACCESSION NUMBER: 2001:921013 CAPLUS
DOCUMENT NUMBER: 136:345886
TITLE: Systematic troubleshooting for LC/MS/MS Part 1:
Sample preparation and chromatography
AUTHOR(S): Naidong, Weng; Halls, Timothy D. J.
CORPORATE SOURCE: Covance Laboratories Inc., Madison, WI, 53704, USA
SOURCE: BioPharm (Duluth, MN, United States) (2001), 14(11), 28,30,32,34,36,38
CODEN: BPRME5; ISSN: 1040-8304
PUBLISHER: Advanstar Communications, Inc.
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylertraline
RL: ANT (Analyte); ANST (Analytical study)
(sample preparation and anal. of drugs in human blood by LC/MS)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS
FORMAT RECORD. ALL CITATIONS AVAILABLE IN THE RE

L4 ANSWER 19 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



L4 ANSWER 19 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
AB Screening methods which enable the selection of selective serotonin reuptake inhibitor compds. which do not possess significant inhibitory potency towards cytochrome P 450 enzymes, in particular, CYP2D6, are disclosed. The invention comprises a method of generating a pharmacophore model of selective serotonin reuptake inhibitor compds. which do not possess significant inhibitory potency towards CYP2D6. Methods for the discovery of selective serotonin reuptake inhibitor compds. which do not possess significant inhibitory potency towards the CYP2D6 enzyme are also disclosed. Pharmaceutical compns. comprising a selective serotonin reuptake inhibitor compound that does not possess significant inhibitory potency towards CYP2D6 as identified by methods of the invention are disclosed. The invention allow the uses of a selective serotonin reuptake inhibitor compound identified by the methods of the invention for the manufacture of medicaments and for the treatment of a condition, a disorder or a disease in a mammal for which such a selective serotonin reuptake inhibitor compound is therapeutically useful.

ACCESSION NUMBER: 2001:691818 CAPLUS
DOCUMENT NUMBER: 135:236399
TITLE: Pharmacophore models for the identification of the CYP2D6 inhibitory potency of selective serotonin reuptake inhibitors
INVENTOR(S): Ekins, Sean
PATENT ASSIGNEE(S): Pfizer Products Inc., USA
SOURCE: Eur. Pat. Appl., 23 pp.
CODEN: EPXXDW
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1134290	A2	20010919	EP 2001-302116	20010308
EP 1134290	A3	20040102		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
US 2002013372	A1	20020131	US 2001-804176	20010312
JP 2001324496	A2	20011122	JP 2001-72613	20010314
PRIORITY APPLN. INFO.:			US 2000-189099P	P 20000314

IT 87857-41-8, Desmethylertraline
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); PRP (Properties); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(pharmacophore models for identification of the CYP2D6 inhibitory potency of selective serotonin reuptake inhibitors)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.


L4 ANSWER 20 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
AB A method of treating asthma that includes the step of controlling the asthma by ingesting a composition which includes a selective serotonin reuptake inhibitor that is sertraline hydrochloride. Chronic administration of the sertraline thereof downregulates brain norepinephrine receptors. The increased output of the brain norepinephrine receptors increases the dilation of the bronchi. Sertraline has no significant affinity for adrenergic (alpha1, alpha2, beta1), cholinergic, GABA, dopaminergic, histaminergic, serotonergic (5HT1A, 5HT1B, 5HT2), or benzodiazepine receptors. Antagonism of such receptors has been hypothesized to be associated with various adverse anticholinergic, sedative, and cardiovascular effects.

ACCESSION NUMBER: 2001:630896 CAPLUS
DOCUMENT NUMBER: 135:175371
TITLE: Composition for treating asthma with a serotonin reuptake inhibitor
INVENTOR(S): Ahmed, Magda Abdel Fattah
PATENT ASSIGNEE(S): USA
SOURCE: U.S., 6 pp.
CODEN: USXXAM
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6281248	B1	20010828	US 2000-505747	20000216
PRIORITY APPLN. INFO.:			US 2000-505747	20000216

IT 87857-41-8, N-Demethylsertraline
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); PROC (Process) (tablets containing sertraline for treating asthma)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 5


THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE

FORMAT

AB Objective: The objective was to evaluate the relationship between the disposition of Sertraline and the presence of the CYP2C19 gene and to define the contribution of cytochrome P 450 2C19 (CYP2C19) to Sertraline N-demethylation. Methods: A single oral 100-mg dose of Sertraline was administered to 6 subjects who were extensive metabolizers and 6 subjects who were poor metabolizers recruited from 77 healthy Chinese volunteers whose genotypes were predicted by polymerase chain reaction-based amplification, followed by restriction fragment length polymorphism analysis. Phenotypes were determined by use of the omeprazole metabolic rate. The

plasma concns. of Sertraline and desmethylsertraline were determined by gas chromatog. with electron-capture detection. Results: Six poor metabolizers with ml mutation had area under the plasma concentration vs. time curve (AUC_{0-∞}) values (983.6 ± 199.3 µg • h/L vs. 697.6 ± 133.0 µg • h/L; P < 0.05) and terminal elimination half-life values of Sertraline (35.5 ± 5.6 h vs. 23.5 ± 4.4 h; P < 0.01) that were significantly higher than the values in 6 extensive metabolizers who were either homozygous or heterozygous for CYP2C19*1. The oral clearance of Sertraline in poor metabolizers (105.3 ± 19.4 L/h) was significantly lower than that of extensive metabolizers (148.4 ± 28.6 L/h). The area under the concentration-time curve from 0 to 144 h and the maximum plasma concentration of desmethylsertraline in poor metabolizers were significantly lower than the values of extensive metabolizers (627.6 ± 203.8 µg • h/L vs. 972.1 ± 270.3 µg • h/L; P < 0.05; and 23.6 ± 6.5 nmol/L vs. 32.4 ± 8.2 nmol/L; P < 0.01; resp.). Conclusions: The polymorphic CYP2C19 appears to be a major enzyme involved in the N-demethylation of Sertraline, and both extensive and poor metabolizers had marked differences in the disposition of Sertraline.

ACCESSION NUMBER: 2001:596898 CAPLUS
DOCUMENT NUMBER: 136:303912
TITLE: Pharmacokinetics of Sertraline in relation to genetic polymorphism of CYP2C19
AUTHOR(S): Wang, Jiu-Hui; Liu, Zhao-Qian; Wang, Wei; Chen, Xiao-Ping; Shu, Yan; He, Nan; Zhou, Hong-Hao
CORPORATE SOURCE: Pharmacogenetics Research Institute, Hunan Medical University, Changsha, 410078, Peop. Rep. China
SOURCE: Clinical Pharmacology & Therapeutics (St. Louis, MO, United States) (2001), 70(1), 42-47
CODEN: CLPTAT; ISSN: 0009-9236
PUBLISHER: Mosby, Inc.
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: BSU (Biological study, unclassified); BIOL (Biological study) (plasma concns. of desmethylsertraline in humans in relation to genetic polymorphism of CYP2C19)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)
Absolute stereochemistry.



REFERENCE COUNT: 25

THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE


FORMAT

AB Recently, clin. depression has been identified as an independent risk factor for increased mortality in patients following acute coronary events. Although the underlying mechanisms of this link remain uncertain, increased platelet activity has been suggested but never proven as the mechanism responsible for this association. Sertraline-HCl is a selective serotonin reuptake inhibitor (SSRI), and is an effective antidepressant agent. Its major liver metabolite, N-demethylsertraline (NDMS), is neurologically inactive. The in vitro effects of escalating concns. of sertraline and NDMS on human platelets were assessed in blood from healthy volunteers and patients with coronary artery disease by aggregometry in plasma and whole blood, by expression of major surface receptors with flow cytometry in washed cells and in the whole blood, and quant. by various platelet function analyzers. Pretreatment of blood samples with sertraline and NDMS resulted in a concentration-dependent inhibition of platelet-rich plasma aggregation induced by 5 µM ADP, by 10 µM ADP, by collagen, and by thrombin. Whole blood platelet aggregability was also reduced when induced by 20 µM ADP and by collagen. Surface expression of surface antigen CD9, glycoprotein (GP) Ib, GP IIb/IIIa, surface antigen VLA-2, P-selectin, and cell adhesion mol. PECAM-1, but not the vitronectin receptor, was also reduced in sertraline- and NDMS-pretreated washed platelets. Whole-blood flow cytometry revealed inhibition of GP IIb/IIIa and P-selectin expression in NDMS-treated samples. Closure time was delayed for the collagen-ADP cartridge and for the collagen-epinephrine cartridge, indicating platelet inhibition in whole blood under high shear conditions. Rapid platelet-function assay revealed a decreased ability of platelets to agglutinate fibrinogen-coated beads, suggesting GP IIb/IIIa inhibition. Thus, both sertraline and its neurologically inactive metabolite, NDMS, exhibited significant concentration-dependent inhibition of human platelets. The antiplatelet effects of sertraline and NDMS may be directly related to the mortality benefits of SSRIs after ischemic events including myocardial infarction and stroke. (c) 2001 The Italian Pharmacological Society.

ACCESSION NUMBER: 2001:411281 CAPLUS
DOCUMENT NUMBER: 136:241479
TITLE: Platelet inhibition by sertraline and N-demethylsertraline: a possible missing link between depression, coronary events, and mortality benefits
of selective serotonin reuptake inhibitors
AUTHOR(S): Serebruany, Victor L.; Gurbel, Paul A.; O'Connor, Christopher M.
CORPORATE SOURCE: Sinai Center for Thrombosis Research, Johns Hopkins University, Baltimore, MD, USA
SOURCE: Pharmacological Research (2001), 43(5), 453-461
CODEN: PHMRP; ISSN: 1043-6618
PUBLISHER: Academic Press
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, N-Demethylsertraline
RL: PAC (Pharmacological activity); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(platelet inhibition by sertraline and desmethylsertraline as link between depression, coronary events, and mortality benefits of

L4 ANSWER 22 OF 79 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
selective serotonin reuptake inhibitors in humans)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 63 THERE ARE 63 CITED REFERENCES AVAILABLE FOR
THIS
FORMAT
RECORD. ALL CITATIONS AVAILABLE IN THE RE


L4 ANSWER 23 OF 79 CAPLUS COPYRIGHT 2004 ACS ON STN
AB The present invention pertains to methods for reducing the platelet
activation state in an individual comprising administering a selective
serotonin reuptake inhibitor (SSRI). The platelet activation state is
reduced upon administering a SSRI, as measured by one or more platelet
activation markers (PAMs). The invention also relates to methods for
treating or preventing an individual at risk for a vascular event,
disease
or disorder by administering a. For example, sertraline-HCl (Zoloft) had
direct platelet inhibitory properties in humans. Moreover,
N-desmethylsertraline, a stable final metabolite of sertraline which was
previously considered inactive, exhibited potent dose-dependent effects
inhibiting human platelets in both platelet rich plasma and in the whole
blood.
ACCESSION NUMBER: 2000:824098 CAPLUS
DOCUMENT NUMBER: 134:522
TITLE: Methods of inhibiting platelet activation with
selective serotonin reuptake inhibitors
INVENTOR(S): Serebruany, Victor L.; Gurbel, Paul A.; O'Connor,
Christopher M.
PATENT ASSIGNEE(S): Heartdrug Research, L.L.C., USA
SOURCE: PCT Int. Appl., 50 pp.
CODEN: PIXXDZ
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000069429	A2	20001123	WO 2000-US13626	20000517
WO 2000069429	A3	20011004		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SI, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
US 6245782	B1	20010612	US 1999-312987	19990517
CA 2375219	AA	20001123	CA 2000-2375219	20000517
EP 1183073	A2	20020306	EP 2000-932556	20000517
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, RO			
AU 766904	B2	20031023	AU 2000-50258	20000517
ZA 2001009994	A	20020826	ZA 2001-9994	20011205
PRIORITY APPLN. INFO.:			US 1999-312987	Al 19990517
			WO 2000-US13626	W 20000517

IT 87857-41-8, N-Demethylsertraline
RL: BAC (Biological activity or effector, except adverse); BSU
(Biological
study, unclassified); BIOL (Biological study)
(inhibition of platelet activation with selective serotonin reuptake
inhibitors for treatment of cardiovascular disorders)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-


L4 ANSWER 23 OF 79 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 24 OF 79 CAPLUS COPYRIGHT 2004 ACS ON STN
AB Three women were studied who were treated during pregnancy with
fluvoxamine, sertraline, or venlafaxine. Amniotic fluid at amniocentesis
and umbilical cord blood and maternal blood at delivery were collected.
and
analyzed for antidepressants and their metabolites by HPLC with UV
detection. The antidepressants and their metabolites were detectable in
all amniotic fluid samples studied, though concns. of the parent compound
were less than those in maternal serum and umbilical cord blood. No
adverse effects of the drugs were reported. The presence of these
antidepressants in amniotic fluid suggests that fetal exposure to these
comps. is continual and may occur through a variety of paths, thus
accounting for increased fetal exposure. These paths include circulatory
via placental passage, gastrointestinal via fetal swallowing, and
respiratory secondary to fetal lung absorption.
ACCESSION NUMBER: 2000:807096 CAPLUS
DOCUMENT NUMBER: 135:28591
TITLE: Amniotic fluid and umbilical cord blood
concentrations
of antidepressants in three women
AUTHOR(S): Hostetter, A.; Ritchie, J. C.; Stowe, Z. N.
CORPORATE SOURCE: Department of Psychiatry and Behavioral Sciences,
Emory University School of Medicine, Atlanta, GA, USA
SOURCE: Biological Psychiatry (2000), 48(10), 1032-1034
CODEN: BIPCBP; ISSN: 0006-3223
PUBLISHER: Elsevier Science Inc.
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Demethylsertraline
RL: BPR (Biological process); BSU (Biological study, unclassified); MFM
(Metabolic formation); BIOL (Biological study); FORM (Formation,
nonpreparative); PROC (Process)
(amniotic fluid and umbilical cord blood concns. of fluvoxamine,
sertraline, venlafaxine, and their metabolites in women)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR
THIS
FORMAT
RECORD. ALL CITATIONS AVAILABLE IN THE RE

L4 ANSWER 25 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB The in vitro biotransformation of bupropion to hydroxybupropion was studied in human liver microsomes and microsomes containing heterologously expressed human cytochromes P 450 (CYP). The mean (\pm S.E.) Km in four human liver microsomes was 89 (\pm 14) μ M. In microsomes containing cDNA-expressed CYPs, hydroxybupropion formation was mediated only by CYP2B6 at 50 μ M bupropion (Km 85 μ M). A CYP2B6 inhibitory antibody produced more than 95% inhibition of bupropion hydroxylation in four human livers. Bupropion hydroxylation activity at 250 μ M was highly correlated with S-mephenytoin N-demethylation activity (yielding nirvanol), another CYP2B6-mediated reaction, in a panel of 32 human livers ($r = 0.94$). The CYP2B6 content of 12 human livers highly correlated with bupropion hydroxylation activity ($r = 0.96$). Thus bupropion hydroxylation is mediated almost exclusively by CYP2B6 and can serve as an index reaction reflecting activity of this isoform. IC50 values for inhibition of a CYP2D6 index reaction (dextromethorphan O-demethylation) by bupropion and hydroxybupropion were 58 and 74 μ M, resp. This suggests a low inhibitory potency vs. CYP2D6, the clin. importance of which is not established. Since bupropion is frequently coadministered with other antidepressants, IC50 values (μ M) for inhibition of bupropion hydroxylation were determined as follows: paroxetine (1.6), fluvoxamine (6.1), sertraline (3.2), desmethylsertraline (19.9), fluoxetine (59.5), norfluoxetine (4.2), and nefazodone (25.4). Bupropion hydroxylation was only weakly inhibited by venlafaxine, O-desmethylvenlafaxine, citalopram, and desmethylcitalopram. The inhibition of bupropion hydroxylation in vitro by a number of newer antidepressants suggests the potential for clin. drug interactions.

ACCESSION NUMBER: 2000:700645 CAPLUS
 DOCUMENT NUMBER: 134:163
 TITLE: CYP2B6 mediates the in vitro hydroxylation of bupropion: potential drug interactions with other antidepressants
 AUTHOR(S): Hesse, Leah M.; Venkatakrishnan, Karthik; Court, Michael H.; Von Moltke, Lisa L.; Duan, Su X.; Shader, Richard I.; Greenblatt, David J.
 CORPORATE SOURCE: Department of Pharmacology and Experimental Therapeutics, New England Medical Center, Tufts University School of Medicine, Boston, MA, 02111, USA
 SOURCE: Drug Metabolism and Disposition (2000), 28(10), 1176-1183
 CODEN: DMSDAI; ISSN: 0090-9556
 PUBLISHER: American Society for Pharmacology and Experimental Therapeutics
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylsertraline
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
 (CYP2B6 mediates in vitro hydroxylation of bupropion: potential drug interactions with other antidepressants)

L4 ANSWER 25 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 41
 THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE
 FORMAT

L4 ANSWER 26 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB Formaldehyde is liberated in the process of cytochrome P 450 (CYP) mediated demethylation of a wide variety of compds. containing the CH3N or CH3O functionality. A highly sensitive method using a high-performance liquid chromatog. (HPLC) system with postcolumn derivatization was developed to measure the liberated formaldehyde as N- and O-demethylase activity of drugs in human liver microsomes. Following the chromatog. separation of formaldehyde on a C18 column, the formaldehyde was reacted with the Nash reagent in the postcolumn reactor at 100° and detected by the fluorescence method. The results showed that the present method has excellent precision and accuracy. The intra- and interassay variances of this method were less than 10%. The newly developed HPLC method was found to be about 80-fold more sensitive than the colorimetric method in detection of formaldehyde. The N-demethylase activity of sertraline in rat liver microsomes determined by the present method did not differ from those detected by previous methods quantifying produced desmethyl metabolite. The present method has been successfully applied to determine the N-demethylase activities of several drugs, including aminopyrine, erythromycin, fluoxetine, S-mephenytoin, and sertraline, in human liver microsomes. This assay should be useful for generic anal. of N- and O-demethylase activities of xenobiotic and endobiotic chems. by CYP enzymes. (c) 2000 Academic Press.

ACCESSION NUMBER: 2000:605323 CAPLUS
 DOCUMENT NUMBER: 133:358817
 TITLE: High-performance liquid chromatography determination of N- and O-demethylase activities of chemicals in human liver microsomes: application of postcolumn fluorescence derivatization using Nash reagent
 AUTHOR(S): Kobayashi, Kaoru; Yamamoto, Tessai; Taguchi, Miwo; Chiba, Kan
 CORPORATE SOURCE: Department of Biochemical Pharmacology and Toxicology, Faculty of Pharmaceutical Sciences, Chiba University, Yayoi-cho, Chiba, 263-8522, Japan
 SOURCE: Analytical Biochemistry (2000), 284(2), 342-347
 CODEN: ANBCA2; ISSN: 0003-2697
 PUBLISHER: Academic Press
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8
 RL: ANF (Analyte); ANST (Analytical study)
 (HPLC determination of N- and O-demethylase activities of chems. in human liver microsomes: postcolumn fluorescence derivatization using Nash reagent)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT:
THIS
FORMAT

14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE

AB This report describes sertraline pharmacokinetics derived from routine therapeutic drug monitoring (TDM) data. A high-performance liquid chromatog. method with UV detection was established for routine sertraline TDM, and 924 analyses were performed from Apr. 1995 to May 1997. Extensive predefined inclusion/exclusion criteria were applied to increase the validity of scientifically evaluated data. Subsequently, 605 samples (65.5%) were excluded. The remaining 319 samples from 319 patients, representative of steady state trough specimens and accompanied by essential clin. information provided on request forms, were scrutinized. A pronounced interindividual variability was observed. Smokers had significantly lower concentration-to-dose (C/D) mean ratios of serum sertraline (s-ert) and its main metabolite desmethylsertraline (s-dsert) than nonsmokers. Higher s-ert and s-dsert C/D mean ratios were found in elderly patients than in adults aged less than 65 yr. In a subset of 20 patients in whom repeated TDM analyses were performed, observed intraindividual sertraline TDM outcome variability was low. The results highlight sertraline TDM as a tool for individual dose optimization and evaluation of patient drug compliance as well as drug-drug interactions.

ACCESSION NUMBER: 2000:601711 CAPLUS
DOCUMENT NUMBER: 134:110069
TITLE: Therapeutic drug monitoring of sertraline:
variability factors as displayed in a clinical setting
AUTHOR(S): Lundmark, Jons; Reis, Margareta; Bengtsson, Finn
CORPORATE SOURCE: Department of Neuroscience and Locomotion, Division of Psychiatry, Linköping University Hospital, Linköping, Swed.
SOURCE: Therapeutic Drug Monitoring (2000), 22(4), 446-454
CODEN: TDMODV; ISSN: 0163-4356
PUBLISHER: Lippincott Williams & Wilkins
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, DESMETHYLSERTRALINE
RL: BPR (Biological process); RSU (Biological study, unclassified); BIOL (Biological study); PROC (Process)
(therapeutic drug monitoring of sertraline)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT:
THIS
FORMAT


35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE

AB Propafenone is mainly metabolized by CYP2D6 to form 5-hydroxypropafenone (5-OHP) and to a minor extent by CYP1A2 and CYP3A4 to form N-depropylpropafenone (N-DPP). The in vitro inhibitory effect of selective serotonin reuptake inhibitors (SSRIs) on the formation of both metabolites was studied, using human liver microsomes. The 5-OHP formation from racemic propafenone and from its individual enantiomers followed 1-enzyme Michaelis-Menten kinetics. Incubation with the racemate yielded a mean V_{max} of 64 pmol \cdot min $^{-1}$ \cdot mg $^{-1}$ and a mean K_m of 0.12 μ M (N = 3). Stereoselectivity in V_{max} and K_m values was observed, with (S)-propafenone displaying higher K_m and V_{max} values. N-DPP formation from racemic propafenone followed 1-enzyme Michaelis-Menten kinetics and yielded a mean V_{max} of 403 pmol \cdot min $^{-1}$ \cdot mg $^{-1}$ and a mean K_m of 116 μ M (N=3). No stereoselectivity in propafenone N-dealkylation was observed. The influence of SSRIs and quinidine, a prototypical CYP2D6 inhibitor, on propafenone 5-hydroxylation was investigated. Quinidine was the most potent inhibitor, followed by fluoxetine, norfluoxetine, and paroxetine. Sertraline, desmethylsertraline, and fluvoxamine had only a moderate inhibitory effect, whereas citalopram displayed slight or no inhibition when racemic propafenone was used as substrate. Mean K_i values of quinidine, fluoxetine, norfluoxetine, and paroxetine were 0.13, 0.33, 0.55, and 0.54 μ M, resp. (N=3). Quinidine and paroxetine were also tested as inhibitors using the individual enantiomers, but no stereoselectivity was observed. Among the SSRIs tested, only fluvoxamine substantially inhibited propafenone N-dealkylation with a mean IC_{50} of 7.0 μ M (N=3). There was a more pronounced inhibitory effect of fluvoxamine on (R)-propafenone than on (S)-propafenone N-dealkylation. In conclusion, these in vitro data suggest that an in vivo interaction between propafenone and the SSRIs, fluoxetine and paroxetine, can be expected, which can lead to clin. relevant β -blockade and an increased risk of side effects in the central nervous system. An interaction with fluvoxamine may be of importance in poor metabolizers for CYP2D6.

ACCESSION NUMBER: 2000:558561 CAPLUS
DOCUMENT NUMBER: 134:95087
TITLE: Effect of selective serotonin reuptake inhibitors on the oxidative metabolism of propafenone. In vitro studies using human liver microsomes
AUTHOR(S): Hemeryck, Alex; De Vriendt, Cindy; Belpaire, Frans M.
CORPORATE SOURCE: Heymans Institute of Pharmacology, Ghent University Medical School, Ghent, 9000, Belg.
SOURCE: Journal of Clinical Psychopharmacology (2000), 20(4), 428-434
CODEN: JCPYDR; ISSN: 0271-0749
PUBLISHER: Lippincott Williams & Wilkins
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: BAC (Biological activity or effector, except adverse); RSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study);
USES (Uses)
(propafenone oxidative metabolism influenced by selective serotonin reuptake inhibitors and possible adverse drug interaction)
RN 87857-41-8 CAPLUS

L4 ANSWER 28 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-
 (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR
 THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE
 FORMAT

L4 ANSWER 29 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB This paper describes a set of simple and sensitive multiresidue methods
 for the determination of the specific serotonin reuptake inhibitors
 (SSRIs) used
 as antidepressant drugs, and some of their resp. active metabolites in
 human serum. It involves liquid-liquid extraction procedures followed
 by gas
 chromatog. coupled to nitrogen phosphorus detection or isocratic
 reversed-phase high-performance liquid chromatog. combined with
 fluorescence
 detection (HPLC-FL), depending on the analytes. Extraction recoveries
 were
 between 71 and 96% for the eight SSRIs and their metabolites analyzed by
 GC and between 41 and 77% for the two of them analyzed by HPLC. Limits
 of
 detection (LODs) and limits of quantitation (LOQs) ranged, resp., from
 2.5
 to 5 µg/l and from 10 to 20 µg/l. Intra-assay and inter-assay
 precision was studied at three and four concentration levels, resp., and
 was less
 than 19% for all compds. Accuracy was also satisfactory for all. An
 excellent linearity was observed from the LOQs up to 1000 µg/l for
 mirtazapine and paroxetine and from each LOQ up to 400 mg/l for the other
 compds. The performance of the methods described thus allows the
 therapeutic drug monitoring of the currently commercialized SSRIs.
 ACCESSION NUMBER: 2000:377581 CAPLUS
 DOCUMENT NUMBER: 133:159587
 TITLE: Methods for the determination of seven selective
 serotonin reuptake inhibitors and three active
 metabolites in human serum using high-performance
 liquid chromatography and gas chromatography
 AUTHOR(S): Lacassie, E.; Gaulier, J.-M.; Marquet, P.; Rabatel,
 J.-F.; Lachatre, G.
 CORPORATE SOURCE: Department of Pharmacology and Toxicology, University
 Hospital, Limoges, 87042, Fr.
 SOURCE: Journal of Chromatography, B: Biomedical Sciences and
 Applications (2000), 742(2), 229-238
 CODEN: JCBREP; ISSN: 0378-4347
 PUBLISHER: Elsevier Science B.V.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylertraline
 RL: BPR (Biological process); BSU (Biological study, unclassified); BIOL
 (Biological study); PROC (Process)
 (methods for determination of seven selective serotonin reuptake
 inhibitors and
 three active metabolites in human serum using high-performance liquid
 chromatog. and gas chromatog.)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-
 (9CI) (CA INDEX NAME)

Absolute stereochemistry.


L4 ANSWER 29 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR
 THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE
 FORMAT

L4 ANSWER 30 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB Grapefruit juice is an inhibitor of cytochrome P 450 3A4 (CYP3A4). This
 study assessed the in vitro and in vivo effects of grapefruit juice on
 sertraline metabolism The in vitro assay involved anal. of sertraline
 metabolism
 by CYP3A4, using CYP3A4-expressed human β -lymphoblast microsomes.
 The in vivo study involved HPLC anal. of serum trough levels of
 sertraline
 and demethylsertraline in patients who had been taking their usual dose
 of
 sertraline for 26 wk, followed by concurrent use of sertraline with
 grapefruit juice for 1 wk. The in vitro assay demonstrated that
 grapefruit juice concentration-dependently inhibited the formation of
 demethylsertraline. In the in vivo study, mean serum sertraline levels
 were determined in patients with a history of depression. The mean
 sertraline
 dosage was 55 mg/day. The results of the in vivo study appeared to be
 consistent with the in vitro findings, in that mean serum sertraline
 trough levels increased from 13.7 µg/L before to 20.2 µg/L after
 administration of grapefruit juice. Thus, the in vitro study
 demonstrated
 that grapefruit juice can inhibit the metabolism of sertraline.
 ACCESSION NUMBER: 2000:29522 CAPLUS
 DOCUMENT NUMBER: 132:58738
 TITLE: The effects of grapefruit juice on sertraline
 metabolism: an in vitro and in vivo study
 AUTHOR(S): Lee, Audrey J.; Chan, William K.; Harralson, Arthur
 F.; Buffum, John; Bul, Bao-Chau Crystal
 CORPORATE SOURCE: School of Pharmacy and Health Sciences, University of
 the Pacific, Stockton, CA, USA
 SOURCE: Clinical Therapeutics (1999), 21(11), 1890-1899
 CODEN: CLTHDG; ISSN: 0149-2918
 PUBLISHER: Excerpta Medica, Inc.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Demethylsertraline
 RL: BPR (Biological process); BSU (Biological study, unclassified); MFM
 (Metabolic formation); BIOL (Biological study); FORM (Formation,
 nonpreparative); PROC (Process)
 (grapefruit juice interaction with sertraline metabolism in humans as
 determined
 by formation of)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-
 (9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR
THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE
FORMAT

AB Aims: The present study was designed to define the kinetic behavior of sertraline N-demethylation in human liver microsomes and to identify the isoforms of cytochrome P 450 involved in this metabolic pathway.

Methods: The kinetics of the formation of N-demethylsertraline were determined in human liver microsomes from six genotyped CYP2C19 extensive (EM) and three poor metabolizers (PM). Selective inhibitors of and specific monoclonal antibodies to various cytochrome P 450 isoforms were also employed. Results: The kinetics of N-demethylsertraline formation in all EM liver microsomes were fitted by a two-enzyme Michaelis-Menten equation, whereas the kinetics in all PM liver microsomes were best described by a single-enzyme Michaelis-Menten equation similar to the low-affinity component found in EM microsomes. Mean apparent Km values for the high- and low-affinity components were 1.9 and 88 µM and Vmax values were 33 and 554 pmol min⁻¹ mg⁻¹ protein, resp., in the EM liver microsomes. Omeprazole (a CYP2C19 substrate) at high concns. and sulfaphenazole (a selective inhibitor of CYP2C9) substantially inhibited N-demethylsertraline formation. Of five monoclonal antibodies to various cytochrome P 450 forms tested, only anti-CYP2C8/9/19 had any inhibitory effect on this reaction. The inhibition of sertraline N-demethylation by anti-CYP2C8/9/19 was greater in EM livers than in PM livers at both low and high substrate concns. However, anti-CYP2C8/9/19 did not abolish the formation of N-demethylsertraline in the microsomes from any of the livers. Conclusions: The polymorphic enzyme CYP2C19 catalyzes the high-affinity N-demethylation of sertraline, while CYP2C9 is one of the low-affinity components of this metabolic pathway.

ACCESSION NUMBER: 1999:653227 CAPLUS
DOCUMENT NUMBER: 132:131735
TITLE: Evidence for involvement of polymorphic CYP2C19 and 2C9 in the N-demethylation of sertraline in human liver microsomes
AUTHOR(S): Xu, Zhen-Hua; Wang, Wei; Zhao, Xue-Jun; Huang, Song-Lin; Zhu, Bing; He, Nan; Shu, Yan; Liu, Zhao-Qian; Zhou, Hong-Hao
CORPORATE SOURCE: Pharmacogenetics Research Institute, Hunan Medical University, Changsha, 410078, Peop. Rep. China
SOURCE: British Journal of Clinical Pharmacology (1999), 48(3), 416-423
CODEN: BCPHBM; ISSN: 0306-5251
PUBLISHER: Blackwell Science Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, N-Demethylsertraline
RL: BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative)
(evidence for involvement of polymorphic CYP2C19 and 2C9 in the N-demethylation of sertraline in human liver microsomes)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR
THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE
FORMAT

AB Three- and four-dimensional quant. structure activity relationship (3D/4D-QSAR) pharmacophore models of competitive inhibitors of CYP2D6 were constructed using data from our laboratory or the literature. The 3D-QSAR pharmacophore models of the common structural features of CYP2D6 inhibitors were built using the program Catalyst (Mol. Simulations, San Diego, CA, USA). These 3D-QSAR models were compared with 3D and 4D-QSAR partial least squares (PLS) models which were constructed using mol. surface-weighted holistic invariant mol. (MS-WHIM) descriptors of size and shape of inhibitors. The first Catalyst model was generated from multiple conformers of competitive inhibitors (n = 20) of CYP2D6 mediated bufuralol 1'-hydroxylation. This model demonstrated a correlation of observed and predicted Ki (apparent) values of r = 0.75. A second Catalyst model was constructed from literature derived Ki (apparent) values (n = 31) for the inhibition of CYP2D6. This model provided a correlation of observed and predicted inhibition for CYP2D6 of r = 0.91. Both Catalyst Ki pharmacophores were then validated by predicting the Ki (apparent) of a test set of known CYP2D6 inhibitors (n = 15). Ten out of 15 of these Ki (apparent) values were predicted to be within one log residual of the observed value using our CYP2D6 inhibitor model, while the literature model predicted nine out of 15 values. Similarly, 3D- and 4D-QSARs derived from PLS MS-WHIM for our dataset yielded predictable models as assessed using cross-validation. The corresponding cross-validated PLS MS-WHIM model for the literature dataset yielded a comparable 3D-QSAR and improved 4D-QSAR value. Such computational models will aid in future prediction of drug-drug interactions.

ACCESSION NUMBER: 1999:635012 CAPLUS
DOCUMENT NUMBER: 132:146140
TITLE: Three and four dimensional-quantitative structure activity relationship (3D/4D-QSAR) analyses of CYP2D6 inhibitors
AUTHOR(S): Ekins, Sean; Bravi, Gianpaolo; Binkley, Shelly; Gillespie, Jennifer S.; Ring, Barbara J.; Wikel, James
CORPORATE SOURCE: H.; Wrighton, Steven A. Departments of Drug Disposition, Lilly Research Laboratories, Eli Lilly and Co., Lilly Corporate Center, Indianapolis, IN, 46285, USA
SOURCE: Pharmacogenetics (1999), 9(4), 477-489
CODEN: PHMCEE; ISSN: 0960-314X
PUBLISHER: Lippincott Williams & Wilkins
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylertraline
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses)
(3D/4D-QSAR analyses of CYP2D6 inhibitors)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-

L4 ANSWER 32 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 49 THERE ARE 49 CITED REFERENCES AVAILABLE FOR
THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE
FORMAT


L4 ANSWER 33 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN

AB Sertraline, a new antidepressant of the selective serotonin re-uptake inhibitor class, is extensively metabolized to desmethylsertraline in humans. We identified the cytochrome P 450 (CYP) isoforms involved in sertraline N-demethylation using pooled human liver microsomes and cDNA-expressed CYP isoforms.adie-Hofstee plots for the sertraline N-demethylation in human liver microsomes were monophasic. The estimated Michaelis-Menten kinetic parameters were: $K_M = 18.1 \pm 2.0 \mu M$, $V_{max} = 0.45 \pm 0.03 \text{ nmol/min/mg of protein}$, and $V_{max}/K_M = 25.2 \pm 4.3 \mu\text{l/min/mg of protein}$. At the substrate concentration of $20 \mu M$, which approximated the apparent K_M value, sulfaphenazole (CYP2C9 inhibitor) and triazolam (CYP3A substrate) reduced the N-demethylation activities by 20 to 35% in human liver microsomes, whereas the inhibition induced by mephenytoin (CYP2C19 substrate) or quinidine (CYP2D6 inhibitor) was marginal. The anti-CYP2B6 antibody inhibited the sertraline N-demethylation activities by 35%. Sertraline N-demethylation activities were detected in all cDNA-expressed CYP isoforms studied. In particular, CYP2C19, CYP2B6, CYP2C9-Arg, CYP2D6-Val, and CYP3A4 all showed relatively high activity. When the contributions of CYP2D6, CYP2C9, CYP2B6, CYP2C19, and CYP3A4 were estimated from the V_{max}/K_M of cDNA-expressed CYP isoforms and from their contents in pooled human liver microsomes, the values were found to be 35, 29, 14, 13, and 9%, resp. The results suggest that at least five isoforms of CYP (CYP2B6, CYP2C9, CYP2C19, CYP2D6, CYP3A4) are involved in the sertraline N-demethylation in human liver microsomes and that the contribution of any individual isoform does not exceed 40% of overall metabolism. Therefore, concurrent administration of a drug that inhibits a specific CYP isoform is unlikely to cause a marked increase in the plasma concentration of sertraline.

ACCESSION NUMBER: 1999:424018 CAPLUS
DOCUMENT NUMBER: 131:193674
TITLE: Sertraline N-demethylation is catalyzed by multiple isoforms of human cytochrome P-450 in vitro
AUTHOR(S): Kobayashi, Kaoru; Ishizuka, Tomoko; Shimada, Noriaki; Yoshimura, Yoshitaka; Kamijima, Kunitoshi; Chiba, Kan
CORPORATE SOURCE: Laboratory of Biochemical Pharmacology and
Toxicology, Faculty of Pharmaceutical Sciences, Chiba University, Chiba, 263-8522, Japan
SOURCE: Drug Metabolism and Disposition (1999), 27(7), 763-766
CODEN: DMDSAI; ISSN: 0090-9556
PUBLISHER: American Society for Pharmacology and Experimental Therapeutics
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: BSU (Biological study, unclassified); MPM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative)
(sertraline N-demethylation is catalyzed by multiple isoforms of human cytochrome P 450 in vitro)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 33 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR
THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE
FORMAT

L4 ANSWER 34 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN

AB (PK) This study was undertaken to determine whether there are pharmacokinetic interactions between zolpidem, a hypnotic, and sertraline, an antidepressant. Twenty-eight healthy female volunteers received a single dose of zolpidem alone and five consecutive dose(s) of zolpidem 10 mg in the presence of chronic doses (19 days) of sertraline 50 mg. Using HPLC, plasma levels of zolpidem, sertraline, and N-desmethylsertraline were determined at different times throughout the study and PK parameters derived. Compared to zolpidem alone, the $T_{1/2}$ of the first dose of zolpidem in the presence of sertraline was reduced, the C_{max} of the fifth zolpidem dose in the presence of sertraline was significantly increased, and its T_{max} was significantly reduced. After five doses of zolpidem, the AUC of sertraline (-6%) and the C_{max} of N-desmethylsertraline (+13%) were changed. There were no next-day effects of zolpidem on the Digit Symbol Substitution Test, and both drugs were well tolerated. Overall, coadministration of sertraline 50 mg and zolpidem 10 mg appears to be safe in healthy females but could result in a shortened onset of action and increased effect of zolpidem.

ACCESSION NUMBER: 1999:339280 CAPLUS
DOCUMENT NUMBER: 131:165239
TITLE: Coadministration of short-term zolpidem with sertraline in healthy women
AUTHOR(S): Allard, Stephane; Sainati, Stephen M.; Roth-Schechter, Barbara F.
CORPORATE SOURCE: Lorex Pharmaceuticals, Skokie, IL, 60680-5110, USA
SOURCE: Journal of Clinical Pharmacology (1999), 39(2), 184-191
CODEN: JPCBR; ISSN: 0091-2700
PUBLISHER: Sage Publications
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8
RL: BPR (Biological process); BSU (Biological study, unclassified); MPM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); PROC (Process)
(drug interactions after coadministration of short-term zolpidem with sertraline in healthy women)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR
THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE
FORMAT

AB The biotransformation of venlafaxine (VF) into its two major metabolites, O-desmethylvenlafaxine (ODV) and N-desmethylvenlafaxine (NDV) was studied in vitro with human liver microsomes and with microsomes containing individual human cytochromes from cDNA-transfected human lymphoblastoid cells. VF was coincubated with selective cytochrome P 450 (CYP) inhibitors and several selective serotonin reuptake inhibitors (SSRIs) to assess their inhibitory effect on VF metabolism. Formation rates for ODV incubated with human microsomes were consistent with Michaelis-Menten kinetics for a single-enzyme mediated reaction with substrate inhibition. Mean parameters determined by non-linear regression were: $V_{max} = 0.36$ nmol/min/mg protein, $K_m = 41 \mu M$, and $K_s 22901 \mu M$ (K_s represents a constant which reflects the degree of substrate inhibition). Quinidine (QUI) was a potent inhibitor of ODV formation with a K_i of $0.04 \mu M$, and paroxetine (PX) was the most potent SSRI at inhibiting ODV formation with a mean K_i value of $0.17 \mu M$. Studies using expressed cytochromes showed that ODV was formed by CYP2C9, -2C19, and -2D6. CYP2D6 was dominant with the lowest K_m , $23.2 \mu M$, and highest intrinsic clearance (V_{max}/K_m ratio). No unique model was applicable to the formation of NDV for all four livers tested. Parameters determined by applying a single-enzyme model were $V_{max} = 2.14$ nmol/min/mg protein, and $K_m = 2504 \mu M$. Ketoconazole was a potent inhibitor of NDV production, although its inhibitory activity was not as great as observed with pure 3A substrates. NDV formation was also reduced by 42% by a polyclonal rabbit antibody against rat liver CYP3A1. Studies using expressed cytochromes showed that NDV was formed by CYP2C9, -2C19, and -3A4. The highest intrinsic clearance was attributable to CYP2C19 and the lowest to CYP3A4. However the high in vivo abundance of 3A isoforms will magnify the importance of this cytochrome. Fluvoxamine (FX), at a concentration of $20 \mu M$, decreased NDV production by 46% consistent with the capacity of FX to inhibit CYP3A, 2C9, and 2C19. These results are consistent with previous studies that show CYP2D6 and -3A4 play important roles in the formation of ODV and NDV, resp. In addition the authors have shown that several other CYPs have important roles in the biotransformation of VF.

ACCESSION NUMBER: 1999:227255 CAPLUS
DOCUMENT NUMBER: 131:67604
TITLE: O- and N-demethylation of venlafaxine in vitro by human liver microsomes and by microsomes from cDNA-transfected cells: effect of metabolic inhibitors and SSRI antidepressants
AUTHOR(S): Fogelman, Steven M.; Schmidt, Jürgen; Venkatakrishnan, Karthik; Von Moltke, Lisa L.; Harmatz, Jerold S.; Shader, Richard I.; Greenblatt, David J.
CORPORATE SOURCE: Department of Pharmacology and Experimental Therapeutics, Tufts University School of Medicine, Boston, MA, 02111, USA
SOURCE: Neuropsychopharmacology (1999), 20(5), 480-490
CODEN: NEROEW; ISSN: 0893-133X
PUBLISHER: Elsevier Science Inc.

DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study) (O- and N-demethylation of venlafaxine in vitro by human liver microsomes and by microsomes from cDNA-transfected cells and role of cytochrome P 450 and effect of metabolic inhibitors and SSRI antidepressants)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR
THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE
FORMAT

AB Objective: To investigate the in vitro potential of selective serotonin reuptake inhibitors (SSRIs) to inhibit two CYP2C9-catalyzed reactions, tolbutamide 4-methylhydroxylation and (S)-warfarin 7-hydroxylation. Methods: The formation of 4-hydroxytolbutamide from tolbutamide and that of 7-hydroxywarfarin from (S)-warfarin as a function of different concns. of SSRIs and some of their metabolites was studied in microsomes from three human livers. Results: Both tolbutamide 4-methylhydroxylation and (S)-warfarin 7-hydroxylation followed one enzyme Michaelis-Menten kinetics. Kinetic anal. of 4-hydroxytolbutamide formation yielded a mean apparent Michaelis-Menten constant (K_m) of $133 \mu M$ and a mean apparent maximal velocity (V_{max}) of $248 \text{ pmol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$; formation of 7-hydroxywarfarin yielded a mean K_m of $3.7 \mu M$ and a mean V_{max} of $10.5 \text{ pmol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$. Amongst the SSRIs and some of their metabolites tested, only fluvoxamine markedly inhibited both reactions. The average computed inhibition constant (K_i) values and ranges of fluvoxamine when tolbutamide and (S)-warfarin were used as substrate, were 13.3 (6.4 - 17.3) μM and 13.0 (8.4 - 18.7) μM , resp. The average K_i value of fluoxetine for (S)-warfarin 7-hydroxylation was 87.0 (57.0 - 125) μM . Conclusion: Amongst the SSRIs tested, fluvoxamine was shown to be the most potent inhibitor of both tolbutamide 4-methylhydroxylation and (S)-warfarin 7-hydroxylation. Fluoxetine, norfluoxetine, paroxetine, sertraline, desmethylsertraline, citalopram, desmethylocitalopram had little or no effect on CYP2C9 activity in vitro. This is consistent with in vivo data indicating that amongst the SSRIs, fluvoxamine has the greatest potential for inhibiting CYP2C9-mediated drug metabolism.

ACCESSION NUMBER: 1999:116414 CAPLUS
DOCUMENT NUMBER: 131:39229
TITLE: Inhibition of CYP2C9 by selective serotonin reuptake inhibitors: in vitro studies with tolbutamide and (S)-warfarin using human liver microsomes
AUTHOR(S): Hemeryck, A.; De Vriendt, C.; Belpaire, F. M.
CORPORATE SOURCE: Heymans Institute of Pharmacology, University of Gent Medical School, Ghent, 9000, Belg.
SOURCE: European Journal of Clinical Pharmacology (1999), 54(12), 947-951
CODEN: EJCPAS; ISSN: 0031-6970
PUBLISHER: Springer-Verlag
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: ADV (Adverse effect, including toxicity); BIOL (Biological study) (serotonin reuptake inhibitors inhibition of CYP2C9-mediated drug metabolism: tolbutamide and (S)-warfarin use as substrates in human liver microsomes)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT:
THIS
FORMAT

28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE

AB A 53-yr-old Caucasian male victim of suicide was suspected of overdose with sertraline and alprazolam after death-scene investigation. The concentration of sertraline, a selective serotonin reuptake inhibitor, was determined by a gas chromatograph with mass selective detection. The concentration of alprazolam, a triazolobenzodiazepine, was determined by high-performance liquid chromatog. The sertraline concentration was reported at 1.0 mg/L in peripheral blood, which is greater than previously reported in other postmortem cases in which death was attributed to a multiple drug overdose. The N-desmethylsertraline concentration was reported at 0.2 mg/L in peripheral blood, which is far less than in other postmortem cases and suggests acute intoxication in this case. The alprazolam concentration was reported at 33 µg/L in heart blood, which is within the therapeutic range. The cause of death was multiple drug intoxication following acute use of sertraline, the manner of death was suicide, and the mechanism of death is an unexplained drug interaction and/or toxicity.

ACCESSION NUMBER: 1998:664934 CAPLUS
DOCUMENT NUMBER: 130:21454
TITLE: Fatal multiple drug intoxication following acute sertraline use
AUTHOR(S): Milner, D. A.; Hall, M.; Davis, G. C.; Brisse, R. M.; Robinson, C. A.
CORPORATE SOURCE: Department of Pathology, Division of Forensic Pathology, University of Alabama at Birmingham, Birmingham, AL, USA
SOURCE: Journal of Analytical Toxicology (1998), 22(6), 545-548
CODEN: JATOD3; ISSN: 0146-4760
PUBLISHER: Preston Publications
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, N-Demethylsertraline
RL: BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative)
(fatal multiple drug intoxication in human following acute sertraline use)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT:
THIS
FORMAT

17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE


AB A veterinary method for clin. modifying the behavior of a household pet dog exhibiting a recognized type of canine affective aggression behavior is provided. The veterinary behavior modification method administers at least one compound selected from the group consisting of R enantiomers, S enantiomers, or a racemic mixture of selective serotonin reuptake inhibitors or their active metabolites to the dog upon one or multiple occasions; and the administration of these compds. will modify clin. the canine affective aggression behavior of the household dog permanently or for an indefinite period of time. This veterinary behavior modification method can be usefully employed as an adjunct to conditioning approaches presently employed and will avoid the need for euthanasia in extreme behavioral circumstances.

ACCESSION NUMBER: 1998:534794 CAPLUS
DOCUMENT NUMBER: 129:156948
TITLE: Modifying the behavior of dogs exhibiting canine affective aggression with R and S enantiomers or racemic mixtures of selective serotonin reuptake inhibitors or their metabolites
INVENTOR(S): Dodman, Nicholas H.
PATENT ASSIGNEE(S): Trustees of Tufts College, USA
SOURCE: U.S., 21 pp., Cont.-in-part of U.S. 5,554,383.
CODEN: USXXAM
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 3
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5788986	A	19980804	US 1996-699112	19960816
US 5554383	A	19960910	US 1995-417747	19950406
PRIORITY APPLN. INFO.:			US 1995-417747	19950406

IT 87857-41-8, Desmethylsertraline
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study);
USES
(Uses)
(modifying the behavior of aggressive dogs with R and S enantiomers or racemic mixts. of selective serotonin reuptake inhibitors or their metabolites)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE
FORMAT

AB A gas chromatog.-mass spectrometry method is presented which allows the simultaneous determination of the plasma concns. of the selective serotonin reuptake inhibitors citalopram, paroxetine, sertraline, and their pharmacol. active N-desmethylated metabolites (desmethylocitalopram, didesmethylcitalopram, desmethylsertraline) after derivatization with the reagent N-methylbis(trifluoroacetamide). No interferences from endogenous compds. are observed following the extraction of plasma samples from six different human subjects. The standard curves are linear over a working range of 10-500 ng/mL for citalopram, 10-300 ng/mL for desmethylocitalopram, 5-60 ng/mL for didesmethylcitalopram, 20-400 ng/mL for sertraline and desmethylsertraline, and 10-200 ng/mL for paroxetine. Recoveries measured at three concns. range from 81 to 188% for the tertiary amines (citalopram and the interla standard methylamprotiline), 73 to 95% for the secondary amines (desmethylocitalopram, paroxetine and sertraline), and 39 to 66% for the primary amines (didesmethylocitalopram and desmethylsertraline). Intra- and interday coeffs. of variation determined at three concns. range from 3 to 11% for citalopram and its metabolites, 4 to 15% for paroxetine, and 5 to 13% for sertraline and desmethylsertraline. The limits of quantitation of the method are 2 ng/mL for citalopram and paroxetine, 1 ng/mL for sertraline, and 0.5 ng/mL for desmethylocitalopram, didesmethylcitalopram, and desmethylsertraline. No interferences are noted from 20 other psychotropic drugs. This sensitive and specific method can be used for single-dose pharmacokinetics. It is also useful for therapeutic drug monitoring of these three drugs and could possibly be adapted for the quantitation of the two other selective serotonin reuptake inhibitors on the market, namely fluoxetine and fluvoxamine.

ACCESSION NUMBER: 1998:458150 CAPLUS
DOCUMENT NUMBER: 129:197500
TITLE: Simultaneous determination of human plasma levels of citalopram, paroxetine, sertraline, and their metabolites by gas chromatography-mass spectrometry
AUTHOR(S): Eap, C. R.; Bouchoux, G.; Amey, M.; Cochar, N.; Savary, L.; Baumann, P.
CORPORATE SOURCE: Unite de Biochimie et Psychopharmacologie Clinique, Departement Universitaire de Psychiatrie Adulte, Prilly-Lausanne, CH-1008, Switz.
SOURCE: Journal of Chromatographic Science (1998), 36(7), 365-371
CODEN: JCHSBE; ISSN: 0021-9665
PUBLISHER: Preston Publications
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: ANT (Analyte); BOC (Biological occurrence); BSU (Biological study, unclassified); MFM (Metabolic formation); ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative); OCCU (Occurrence) (simultaneous determination of human plasma levels of citalopram, paroxetine,

L4 ANSWER 39 OF 79 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE
FORMAT

AB To characterize milk/plasma (M/P) ratio and infant exposure, for sertraline and N-desmethylsertraline, in breast-feeding women taking sertraline for the treatment of depression. Eight women (mean age 28 yr) taking sertraline (1.05 mg kg⁻¹ day⁻¹) and their infants (mean age 5.7 mo) were studied. Sertraline and N-desmethylsertraline in plasma and milk were measured by high-performance liquid chromatog. over a 24 h dose interval at steady-state. M/P values were estimated from area under the plasma and milk concentration-time curves. All milk produced was collected over the dose interval. Infant exposure was estimated as the product of actual or estimated milk production, and average drug concentration in milk, normalized to body weight and expressed as a percentage of the weight-adjusted maternal dose. Mean milk production was 321 mL day⁻¹ (range 34-974 mL). Mean M/P values of 1.93 and 1.64 were calculated for sertraline and N-desmethylsertraline resp. Infant exposure estimated from actual milk produced was 0.2% and 0.3% of the weight-adjusted maternal dose for sertraline and N-desmethylsertraline (as sertraline equivalent) resp. When calculated from estimated milk production (0.15 l kg⁻¹ day⁻¹), infant exposure was significantly greater (P<0.0001) at 0.90% and 1.32% for sertraline and N-desmethylsertraline resp. Neither sertraline nor its N-desmethyl metabolite could be detected in plasma samples from the four infants tested. No adverse effects were observed in any of the eight infants and all had achieved normal developmental milestones. Irresp. of the method of calcn. of infant exposure, the mean total dose of sertraline and its N-desmethyl metabolite transmitted to infants via breast-feeding is low and unlikely to cause any significant adverse effects.

ACCESSION NUMBER: 1998:375347 CAPLUS
DOCUMENT NUMBER: 129:156408
TITLE: Distribution and excretion of sertraline and N-desmethylsertraline in human milk
AUTHOR(S): Kristensen, J. H.; Ilett, K. F.; Ducei, L. J.; Hackett, L. P.; Yapp, P.; Wojnar-Horton, R. E.; Roberts, M. J.; Paech, M.
CORPORATE SOURCE: Department of Pharmacy, King Edward Memorial and Princess Margaret Hospitals, Subiaco, 6008, Australia
SOURCE: British Journal of Clinical Pharmacology (1998), 45(5), 453-457
CODEN: BCPHBM; ISSN: 0306-5251
PUBLISHER: Blackwell Science Ltd.
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8
RL: BPR (Biological process); BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); PROC (Process) (distribution and excretion of sertraline and N-desmethylsertraline in human milk)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT:
THIS
FORMAT


23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE

AB Objective: Biotransformation of metoprolol to α -hydroxymetoprolol (HM) and O-demethylmetoprolol (ODM) is mediated by CYP2D6. The selective serotonin reuptake inhibitors (SSRIs) are known to inhibit CYP2D6. The aim was to study in vitro the potential inhibitory effect of SSRIs on metoprolol biotransformation. Methods: Using microsomes from two human livers, biotransformation of metoprolol to α -hydroxymetoprolol (HM) and O-demethylmetoprolol (ODM) as a function of the concns. of the SSRIs and of some of their metabolites was studied. Results: The kinetics of the formation of both metabolites are best described by a biphasic enzyme model. The estimated values of Vmax and KM for the high affinity site

are for the α -hydroxylation in human liver HL-1 32 pmol mg⁻¹ min⁻¹ and 75 μ mol \cdot l⁻¹ resp., and in human liver HL-9 39 pmol mg⁻¹ min⁻¹ and 70 μ mol \cdot l⁻¹ resp.; for the O-demethylation in HL-1 131 pmol mg⁻¹ min⁻¹ and 95 μ mol \cdot l⁻¹ resp., and in HL-9 145 pmol mg⁻¹ min⁻¹ and 94 μ mol \cdot l⁻¹ resp. Quinidine is for both pathways a potent inhibitor of the high-affinity site, with Ki values ranging from 0.03 to 0.18 μ mol \cdot l⁻¹. Fluoxetine, norfluoxetine and paroxetine are likewise potent inhibitors, with Ki values ranging from 0.30 to 2.1 μ mol \cdot l⁻¹ fluvoxamine, sertraline, desmethylsertraline, citalopram and desmethylcitalopram are less potent inhibitors, with Ki values above 10 μ mol \cdot l⁻¹. Conclusion: The rank order of the SSRIs for inhibition of metoprolol metabolism is comparable to that reported in the literature for other CYP2D6 substrates, with fluoxetine, norfluoxetine and paroxetine being the most potent. These findings need further investigation to determine their clin. relevance.

ACCESSION NUMBER: 1998:327690 CAPLUS
DOCUMENT NUMBER: 129:156445
TITLE: The oxidative metabolism of metoprolol in human liver microsomes: inhibition by the selective serotonin reuptake inhibitors
AUTHOR(S): Belpaire, F. M.; Wijnant, P.; Temmermann, A.; Rasmussen, B. B.; Brusen, K.
CORPORATE SOURCE: Heymans Institute of Pharmacology, University of Gent Medical School, De Pintelaan 185, Ghent, B-9000, Belg.
SOURCE: European Journal of Clinical Pharmacology (1998), 54(3), 261-264
CODEN: EJCPAS; ISSN: 0031-6970
PUBLISHER: Springer-Verlag
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study) (oxidative metabolism of metoprolol in human liver microsomes and inhibition by selective serotonin reuptake inhibitors)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT:
THIS
FORMAT


22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE

AB The extent of changes in CYP2D6 and CYP1A2 activities with higher therapeutic dosages (>50 mg/day) of sertraline is not well established in vivo. This study assessed the extent and determinants of changes in CYP2D6 and CYP1A2 isoenzyme activities after treatment with clin.

relevant doses of sertraline. Patients and healthy volunteers aged 19 to 85 yr (N = 21) were treated with sertraline for 5 to 35 days. The dosage of sertraline ranged from 25 to 150 mg/day (93.3 \pm 26.4 mg/day; mean \pm SD). All subjects had an extensive metabolizer phenotype for CYP2D6 and received a single oral dose of dextromethorphan (30 mg) and caffeine (100 mg) before and after sertraline treatment. The log O-demethylation ratio (ODMR) of dextromethorphan and the caffeine metabolic ratio (CMR) in overnight urine were used as in vivo indexes of the CYP2D6 and CYP1A2 isoenzyme activities, resp. Concurrent medications and lifestyle habits (e.g., smoking and diet) were monitored during the study. Baseline log ODMR (-2.33 \pm 0.45) but not CMR (5.1 \pm 1.9) (mean \pm SD) significantly changed after sertraline treatment (-2.19 \pm 0.62; 4.5 \pm 1.6, resp.) (p: ODMR = 0.04, CMR = 0.10). There was no significant effect of age, dose, duration of treatment, gender, sertraline and/or desmethylsertraline plasma concentration, subject type (patient or volunteer), and weight on the extent of changes in log ODMR or CMR (p > 0.05). In conclusion, sertraline treatment at a mean daily dosage of 94.0 mg did not significantly change CYP1A2 activity and resulted in a modest inhibition of CYP2D6 activity.

ACCESSION NUMBER: 1998:88540 CAPLUS
DOCUMENT NUMBER: 128:212643
TITLE: The extent and determinants of changes in CYP2D6 and CYP1A2 activities with therapeutic doses of sertraline
AUTHOR(S): Ozdemir, Vural; Naranjo, Claudio A.; Herrmann, Nathan; Shulman, Richard W.; Sellers, Edward M.; Reed, Ken; Kalow, Werner
CORPORATE SOURCE: Sunnybrook Health Science Centre, Women's College Hospital, Toronto, ON, Can.
SOURCE: Journal of Clinical Psychopharmacology (1998), 18(1), 55-61
CODEN: JCPYDR; ISSN: 0271-0749
PUBLISHER: Williams & Wilkins
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: BOC (Biological occurrence); BSU (Biological study, unclassified); MFH (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); OCCU (Occurrence) (sertraline effects on CYP2D6 and CYP1A2 activities in humans)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT:
THIS
FORMAT

33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE

AB Several new antidepressants that inhibit the serotonin (SERT) and norepinephrine transporters (NET) have been introduced into clin. practice the past several years. This report focuses on the further pharmacol. characterization of nefazodone and its metabolites within the serotonergic and noradrenergic systems, in comparison with other antidepressants. By use of radioligand binding assays, we measured the affinity (K_i) of 13 antidepressants and 6 metabolites for the rat and human SERT and NET. The K_i values for eight of the antidepressants and three metabolites were also determined for the rat 5-HT_{1A}, 5-HT_{2A} and muscarinic cholinergic receptors, the guinea pig histamine₁ receptor and the human alpha-1 and alpha-2 receptors. These data are useful for predicting side effect profiles and the potential for pharmacodynamic drug-drug interactions of antidepressants. Of particular interest were the findings that paroxetine, generally thought of as a selective SERT antagonist, possesses moderately high affinity for the NET and that venlafaxine, which has been described as a "dual uptake inhibitor", possesses weak affinity for the NET. We observed significant correlations in SERT (r = 0.965) or NET (r = 0.983) affinity between rat and human transporters. Significant correlations were also observed between muscarinic cholinergic and NET affinity. There are several significant correlations between affinities for the 5-HT_{1A}, 5-HT_{2A}, histamine₁, alpha-1 and alpha-2 receptors. These novel findings, not widely described previously, suggest that many of the individual drugs studied in these expts. possess some structural characteristic that detes. affinity for several G protein-coupled, but not muscarinic, receptors.

ACCESSION NUMBER: 1998:26557 CAPLUS
DOCUMENT NUMBER: 128:162774
TITLE: Neurotransmitter receptor and transporter binding profile of antidepressants and their metabolites
AUTHOR(S): Owens, Michael J.; Morgan, W. Neal; Plott, Susan J.; Nemeroff, Charles B.
CORPORATE SOURCE: Laboratory of Neuropsychopharmacology, Department of Psychiatry and Behavioral Sciences, Emory University School of Medicine, Atlanta, Georgia
SOURCE: Journal of Pharmacology and Experimental Therapeutics (1997), 283(3), 1305-1322
CODEN: JPETAB; ISSN: 0022-3565
PUBLISHER: Williams & Wilkins
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study) (binding profile of antidepressants and their metabolites to serotonin and norepinephrine receptors and transporters)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT:
THIS
FORMAT

43 THERE ARE 43 CITED REFERENCES AVAILABLE FOR
RECORD. ALL CITATIONS AVAILABLE IN THE RE

AB Using radioligand binding assays, we determined the equilibrium dissociation consts. (K_D's) for 37 antidepressants, three of their metabolites (desmethylcitalopram, desmethylsertraline, and norfluoxetine), some mood stabilizers, and assorted other compds. (some antiepileptics, Ca²⁺ channel antagonists, benzodiazepines, psychostimulants, antihistamines, and monoamines) for the human serotonin, norepinephrine, and dopamine transporters. Among the compds. that we tested, mazindol was the most potent at the human norepinephrine and dopamine transporters with K_D's of 0.45±0.03 nM and 8.1±0.4 nM, resp. Sertraline (K_D=25±2 nM) and nomifensine (56±3 nM) were the two most potent antidepressants at the human dopamine transporter. We showed significant correlations for antidepressant affinities at binding to serotonin (R=0.93), norepinephrine (R=0.97), and dopamine (R=0.87) transporters in comparison to their resp. values for inhibiting uptake of monoamines into rat brain synaptosomes. These data are useful in predicting some possible adverse effects and drug-drug interactions of antidepressants and related compds.

ACCESSION NUMBER: 1997:808034 CAPLUS
DOCUMENT NUMBER: 128:149531
TITLE: Pharmacological profile of antidepressants and related compounds at human monoamine transporters
AUTHOR(S): Tatsumi, Masahiko; Groshan, Karen; Blakely, Randy D.; Richelson, Elliott
CORPORATE SOURCE: San Pablo Road, Mayo Clinic Jacksonville, Jacksonville, FL 32224, 4500, USA
SOURCE: European Journal of Pharmacology (1997), 340(2/3), 249-258
CODEN: EJPHAZ; ISSN: 0014-2999
PUBLISHER: Elsevier Science B.V.
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: ADV (Adverse effect, including toxicity); BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); THU (Therapeutic use); BIOL (Biological study); USES (Uses) (pharmacol. profile of antidepressants and related compds. at human monoamine transporters)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L4 ANSWER 45 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB A HPLC method for sertraline and desmethylsertraline determination in human plasma and red blood cells is described.
 ACCESSION NUMBER: 1997:779112 CAPLUS
 DOCUMENT NUMBER: 128:70336
 TITLE: Analysis of sertraline and desmethylsertraline in human plasma and red blood cells
 AUTHOR(S): Vataassery, Govind T.; Holden, Lori A.; Hazel, Dana K.; Dysken, Maurice W.
 CORPORATE SOURCE: VA Medical Center, GRECC Program, Research Service, Minneapolis, MN, 55417, USA
 SOURCE: Clinical Biochemistry (1997), 30(7), 565-568
 CODEN: CLBIAS; ISSN: 0009-9120
 PUBLISHER: Elsevier Science Inc.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylsertraline
 RL: ANT (Analyte); ANST (Analytical study)
 (sertraline and desmethylsertraline determination in human plasma and red blood cells by HPLC)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)


Absolute stereochemistry.



REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L4 ANSWER 46 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB Inhibition of cytochrome P 450 (CYP) activity by selective serotonin reuptake inhibitors (SSRIs) has frequently been reported with regard to pathways mediated by CYP2D6, CYP3A4/5, and CYP1A2. Little data exist on the capability of SSRIs to inhibit CYP2C9. We investigated the effect of SSRIs on p-hydroxylation of phenytoin (PH), an established index reaction reflecting CYP2C9 activity, in an in vitro assay using liver tissue from six different human donors. In control incubations (without inhibitor), 5-(p-hydroxy-phenyl)-5-phenylhydantoin (HPH) formation rates were: Vmax 0.023 nmol min⁻¹ mg⁻¹; Km 14.3 μM. Average inhibition consts. (Ki) differed significantly among the SSRIs, with fluvoxamine having the lowest Ki (6 μM) followed by R-fluoxetine (13 μM), norfluoxetine (17 μM), RS-fluoxetine (19 μM), sertraline (33 μM), paroxetine (35 μM), S-fluoxetine (62 μM), and desmethylsertraline (66 μM). Thus, assuming comparable molar concns. at the site of inhibition, fluvoxamine can be expected to have the highest probability of interfering with the metabolism of CYP2C9 substrates. S-fluoxetine is on average a 5 fold weaker CYP2C9 inhibitor, than either R-fluoxetine or the racemic mixture. These findings are consistent with published case reports describing SSRI-related increments in plasma phenytoin levels. Because phenytoin has a narrow therapeutic index, plasma levels should be closely monitored when SSRIs are coadministered.
 ACCESSION NUMBER: 1997:752411 CAPLUS
 DOCUMENT NUMBER: 128:84323
 TITLE: Inhibition of CYP2C9 by selective serotonin reuptake inhibitors in vitro: studies of phenytoin p-hydroxylation
 AUTHOR(S): Schmider, Jurgen; Greenblatt, David J.; Von Moltke, Lisa L.; Karsov, Dmitry; Shader, Richard I.
 CORPORATE SOURCE: Department of Pharmacology and Experimental Therapeutics, Tufts University School of Medicine, Boston, MA, 02111, USA
 SOURCE: British Journal of Clinical Pharmacology (1997), 44(5), 495-498
 CODEN: BCPHEM; ISSN: 0306-5251
 PUBLISHER: Blackwell Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylsertraline
 RL: ADV (Adverse effect, including toxicity); BIOL (Biological study)
 (inhibition of CYP2C9 by selective serotonin reuptake inhibitors)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L4 ANSWER 47 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB Biotransformation of the H-1 antagonist terfenadine to its desalkyl and hydroxy metabolites was studied in vitro using microsomal preps. of human liver. These metabolic reactions are presumed to be mediated by Cytochrome P 450-3A isoforms. The azole antifungal agent ketoconazole was a highly potent inhibitor of both reactions, having mean inhibition consts. (K_i) of 0.037 and 0.34 μ M for desalkyl- and hydroxy-terfenadine formation, resp. Itraconazole also was a potent inhibitor, with K_i values of 0.28 and 2.05 μ M, resp. Fluconazole, on the other hand, was a weak inhibitor. Six selective serotonin reuptake inhibitor antidepressants tested in this system were at least 20 times less potent inhibitors of terfenadine metabolism than was ketoconazole. An in vitro-in vivo scaling model used in vitro K_i values, typical clin. relevant plasma concns. of inhibitors, and presumed liver:plasma partition ratios to predict the degree of terfenadine clearance impairment during coadministration of terfenadine with these inhibitors in humans. The model predicted a large and potentially hazardous impairment of terfenadine clearance by ketoconazole and, to a slightly lesser extent, by itraconazole. However, fluconazole and the six selective serotonin reuptake inhibitors (SSRIs) at usual clin. doses were not predicted to impair terfenadine clearance to a degree that would be of clin. importance. Caution is nonetheless warranted with the coadministration of SSRIs and terfenadine when high doses of SSRIs (particularly fluoxetine) are administered. Also, some individuals may be unusually susceptible to metabolic inhibition for a variety of reasons.

ACCESSION NUMBER: 1997:702816 CAPLUS
 DOCUMENT NUMBER: 128:10069
 TITLE: Inhibition of terfenadine metabolism in vitro by azole antifungal agents and by selective serotonin reuptake inhibitor antidepressants: relation to pharmacokinetic interactions in vivo
 AUTHOR(S): von Moltke, Lisa L.; Greenblatt, David J.; Duan, Su Xiang; Harmatz, Jerold S.; Wright, Eugene C.; Shader, Richard I.
 CORPORATE SOURCE: Dep. of Pharmacol. and Exp. Therapeutics, Div. of Clin. Pharmacol., New England med. Cent. Hosp., Tufts Univ. Sch. of Med., Boston, MA, 02111, USA
 SOURCE: Journal of Clinical Psychopharmacology (1996), 16(2), 104-112
 PUBLISHER: Williams & Wilkins
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylsertraline
 RL: ADV (Adverse effect, including toxicity); BIOL (Biological study) (antifungal and antidepressants drugs interaction with terfenadine metabolism and pharmacokinetics)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

L4 ANSWER 47 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



REFERENCE COUNT: 51 THERE ARE 51 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT


Absolute stereochemistry.

L4 ANSWER 48 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB A HPLC method was developed for determination of risperidone and its therapeutically active main metabolite 9-hydroxyrisperidone in serum. After a single-step liquid-liquid extraction the analytes were separated on a C18 column and measured by UV detection at 280 nm. Inter-day coefficient of variation was <7% for both compds. at serum levels occurring in patients treated with ordinary doses. Studies of anal. interference showed that the most commonly coadministered antidepressants and benzodiazepines did not interfere. Some conventional low dose neuroleptics and clozapine did interfere, but this is of minor importance, because risperidone is intended as an alternative to these drugs.

ACCESSION NUMBER: 1997:573519 CAPLUS
 DOCUMENT NUMBER: 127:272141
 TITLE: Simplified high-performance liquid chromatographic method for determination of risperidone and 9-hydroxyrisperidone in serum from patients comedicated with other psychotropic drugs
 AUTHOR(S): Olesen, Ole Vendelin; Linnet, Kristian
 CORPORATE SOURCE: Institute for Basic Research in Psychiatry, Department of Biological Psychiatry, Clinical Biochemical Laboratory, Psychiatric Hospital, Aarhus University Hospital, Risskov, DK-8240, Den.
 SOURCE: Journal of Chromatography, B: Biomedical Sciences and Applications (1997), 698(1 + 2), 209-216
 CODEN: JCBBEF; ISSN: 0378-4347
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Demethylsertraline
 RL: ADV (Adverse effect, including toxicity); BIOL (Biological study) (HPLC assay for risperidone and 9-hydroxyrisperidone determination in serum from patients comedicated with other psychotropic drugs)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

L4 ANSWER 48 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 REFORMAT RECORD. ALL CITATIONS AVAILABLE IN THE RE

Absolute stereochemistry.



REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS


L4 ANSWER 49 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN

AB A nonblinded study was conducted to compare the pharmacokinetic properties of the selective serotonin reuptake inhibitor sertraline in 22 young (aged 18 to 45 yr) and 22 elderly (> 65 yr) volunteers, of whom half were male and half were female. In this study, sertraline was administered at a dosage of 200mg once daily (the maximum recommended daily dosage) for 21 days after upward dosage titration from 50 mg/day over a 9-day period. Thus, this study was designed to measure the effect of age and gender on the pharmacokinetic properties of sertraline at the maximum dosage recommended for clin. use. The terminal elimination half-life (t_{1/2β}) of sertraline was similar in young females, elderly males and elderly females (mean t_{1/2β} ranged from 32.1 to 36.7 h in these groups) but shorter (22.4 h) in the young males. The mean maximum plasma sertraline concentration (C_{max}) and the mean steady-state area under the plasma concentration-time curve from time zero to 24 h postdose (AUC₂₄) were also similar between the young females, elderly males and elderly females, but were approx. 25% lower in the young males. The time to C_{max} was unaffected by age or gender and ranged from 6.4 to 6.9 h. N-Demethylsertraline is the principal metabolite of sertraline and does not contribute significantly to its serotonergic actions. The mean values for N-demethylsertraline trough plasma concns., AUC₂₄ and C_{max} were comparable in elderly males and females and young females but lower in young males. The ratios of mean AUC₂₄ and C_{max} for N-demethylsertraline to the AUC₂₄ and C_{max} for sertraline were similar between the 4 groups.

ACCESSION NUMBER: 1997:212626 CAPLUS
DOCUMENT NUMBER: 126:246336
TITLE: Pharmacokinetics of sertraline and its N-demethyl metabolite in elderly and young male and female volunteers
AUTHOR(S): Ronfeld, Robert A.; Tremaine, Larry M.; Wilner, Keith D.
CORPORATE SOURCE: Pfizer Central Research, Groton, CT, USA
SOURCE: Clinical Pharmacokinetics (1997), 32(Suppl. 1), 22-30
CODEN: CPKNDH; ISSN: 0312-5963
PUBLISHER: Adis
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87957-41-8, N-Demethylsertraline
RL: BPR (Biological process); BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); PROC (Process)
(pharmacokinetics of sertraline and its N-demethyl metabolite in elderly and young male and female volunteers)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 49 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




L4 ANSWER 50 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN

AB Initial expts. demonstrated that the original CEDIA (cloned enzyme donor immunoassay) benzodiazepine assay cross-reacted with sertraline and sertraline metabolites. In response to this phenomenon, Boehringer Mannheim Corporation developed an improved CEDIA benzodiazepine assay in order to eliminate sertraline cross-reactivity. The improved CEDIA assay was evaluated against the original CEDIA product, EMIT II (enzyme multiplied immunoassay technique) benzodiazepine assay and electron capture neg. chemical ionization (ECNCI) gas chromatog.-mass spectrometry (GC-MS). Five hundred and thirty-one urine drug screens were tested by the immunoassays. Sensitivity and specificity of these immunoassays for the 5-aryl-7-chloro-1,4-benzodiazepine compds. were 92 and 98%, resp.,

for the improved CEDIA assay: 92 and 93%, resp., for the current CEDIA assay; and 87 and 98%, resp., for EMIT II. The improved CEDIA assay performed almost identically to the EMIT II assay, both of which had a significant advantage over the origin CEDIA product, which was subject to cross-reactivity because of sertraline metabolites. The α-hydroxyketone metabolites of sertraline are identified in human urine specimens for the first time using ECNCI GC-MS.

ACCESSION NUMBER: 1997:77332 CAPLUS
DOCUMENT NUMBER: 126:196148
TITLE: Improved CEDIA benzodiazepine assay eliminates sertraline cross-reactivity
AUTHOR(S): Fitzgerald, Robert L.; Herold, David A.
CORPORATE SOURCE: Veterans Affairs Medical Center, Univ. California, San Diego, CA, 92161, USA
SOURCE: Journal of Analytical Toxicology (1997), 21(1), 32-35
CODEN: JATOD3; ISSN: 0146-4760
PUBLISHER: Preston Publications
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8
RL: ANVT (Analyte); ANST (Analytical study)
(Improved CEDIA benzodiazepine assay for elimination of sertraline cross-reactivity in human urine)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 51 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB Biotransformation of phenacetin via O-de-ethylation to acetaminophen, an index reaction reflecting activity of Cytochrome P 450-1A2, was studied in microsomal preps. from a series of human livers. Acetaminophen formation was consistent with a double Michaelis-Menten system, with low-Km (mean $K_{m1} = 68 \mu\text{M}$) and high-Km (mean $K_{m2} = 7691 \mu\text{M}$) components. The low-Km enzyme accounted for an average of 96% of estimated intrinsic clearance, and was predicted to contribute more than 50% of net reaction velocity at phenacetin concns. less than $2000 \mu\text{M}$. Among index inhibitor probes, α -naphthoflavone was a highly potent inhibitor of the low-Km enzyme ($K_{i1} = 0.013 \mu\text{M}$); furafylline also was a moderately active inhibitor ($K_{i1} = 4.4 \mu\text{M}$), but its inhibiting potency was increased by preincubation with microsomes. Ketocozazole was a relatively weak inhibitor ($K_{i1} = 32 \mu\text{M}$); quinidine and cimetidine showed minimal inhibiting activity. Among six selective serotonin reuptake inhibitor (SSRI) antidepressants, fluvoxamine was a potent inhibitor of 1A2 (mean $K_{i1} = 0.24 \mu\text{M}$). The other SSRIs were more than ten-fold less potent. Mean K_{i1} values were: fluoxetine, $4.4 \mu\text{M}$; norfluoxetine, $15.9 \mu\text{M}$; sertraline, $8.8 \mu\text{M}$; desmethylsertraline, $9.5 \mu\text{M}$; paroxetine, $5.5 \mu\text{M}$. The antidepressant nefazodone and four of its metabolites (meta-chloro-phenylpiperazine, two hydroxylated derivs., and a triazolidione) were very weak inhibitors of P 450-1A2. Venlafaxine and its O- and N-desmethyl metabolites showed minimal inhibitory activity.

ACCESSION NUMBER: 1997:66943 CAPLUS
 DOCUMENT NUMBER: 126:246342
 TITLE: Phenacetin O-deethylation by human liver microsomes in vitro. Inhibition by chemical probes, SSRI antidepressants, nefazodone and venlafaxine


AUTHOR(S): Von Moltke, Lisa L.; Greenblatt, David J.; Duan, Su Xiang; Schmider, Juergen; Kudchadker, Leena; Fogelmann, Steven M.; Harmatz, Jerold S.; Shader, Richard I.
 CORPORATE SOURCE: School Medicine, Tufts University, Boston, MA, 02111, USA
 SOURCE: Psychopharmacology (Berlin) (1996), 128(4), 398-407
 CODEN: PSCHDL; ISSN: 0033-3158
 PUBLISHER: Springer
 DOCUMENT TYPE: Journal
 LANGUAGE: English

IT 87857-41-8, Desmethylsertraline
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)

(phenacetin O-deethylation by human liver microsomes in vitro, inhibition by chemical probes, SSRI antidepressants, nefazodone, and venlafaxine)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)


Absolute stereochemistry.

L4 ANSWER 51 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



L4 ANSWER 52 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 52 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB Substrates and inhibitors of the cytochrome P 450 isoenzyme CYP2D6 have overlapping structural characteristics. Two prototype serotonin uptake inhibitors, sertraline and fluoxetine, share these structural criteria and have been identified as potent inhibitors of CYP2D6 in vitro. The current study was undertaken to investigate whether genetically determined CYP2D6 activity alters the disposition of sertraline or fluoxetine or both. Single doses of sertraline (50 mg) and fluoxetine (20 mg) were administered successively to 20 young men with high (extensive metabolizers; n = 10) and low (poor metabolizers; n = 10) CYP2D6 activity. Blood and urine samples were collected for 5 to 7 half-lives and sertraline, desmethylsertraline, fluoxetine, and norfluoxetine were determined by GC and HPLC techniques. Poor metabolizers had significantly greater fluoxetine peak plasma concns. (C_{max} ; 57%), area under the concentration vs. time curve (AUC_{0-∞}; 290%), and terminal elimination half-life (216%) compared with extensive metabolizers. The total amount of fluoxetine excreted in the urine during 8 days was almost three times higher in poor metabolizers than in extensive metabolizers (719 vs. 225 μg ; $p < 0.05$), whereas the total amount of norfluoxetine excreted in urine of poor metabolizers was about half of that of extensive metabolizers (524 vs. 1047 μg ; $p < 0.05$). Norfluoxetine C_{max} and AUC_{0-∞} were significantly smaller in poor metabolizers (55% and 53%, resp.), and the partial metabolic clearance of fluoxetine into norfluoxetine was 10 times smaller in this group (4.3 ± 1.9 vs. 0.4 ± 0.1 L/h; $p < 0.05$). No significant differences between extensive and poor metabolizers were found for sertraline and desmethylsertraline pharmacokinetics. These data indicate that poor metabolizers accumulate fluoxetine but not sertraline and that CYP2D6 plays an important role in the demethylation of fluoxetine but not of sertraline.

ACCESSION NUMBER: 1997:8449 CAPLUS
 DOCUMENT NUMBER: 126:42232
 TITLE: The disposition of fluoxetine but not sertraline is altered in poor metabolizers of debrisoquin

AUTHOR(S): Hamelin, Bettina A.; Turgeon, Jacques; Vallee, Francois; Belanger, Pierre-Maxime; Paquet, Francois; LeBel, Marc
 CORPORATE SOURCE: School of Pharmacy, the Quebec Heart Institute, Laval Hospital, and Anapharm Inc., Universite Laval, Sainte-Foy, QC, Can.

SOURCE: Clinical Pharmacology and Therapeutics (St. Louis) (1996), 60(5), 512-521
 CODEN: CLPTAT; ISSN: 0009-9236
 PUBLISHER: Mosby-Year Book
 DOCUMENT TYPE: Journal
 LANGUAGE: English

IT 87857-41-8, Desmethylsertraline
 RL: BPR (Biological process); BSU (Biological study, unclassified); MFH (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); PROC (Process)


(disposition of fluoxetine but not sertraline is altered in poor metabolizers of debrisoquin)
 RN 87857-41-8 CAPLUS

Absolute stereochemistry.

L4 ANSWER 53 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB This report introduces a flow-based extraction method where an aqueous sample and organic solvent are injected sequentially into an extraction coil, then mixed and separated due to the differential flow velocities of the aqueous and organic phases. A 500 µl aqueous sample is propelled through a 50 µl segment of organic solvent whose flow is impeded due to hydrophobic interactions with the walls of a Teflon extraction coil. This wall drag allows the faster moving aqueous sample to penetrate through and ultimately sep. from the slower organic solvent. These steps are repeated with a back extraction into a second aqueous segment (100µl) that is collected and analyzed with high-pressure liquid chromatog. (HPLC). The configuration of this novel sequential-injection extraction (SIE) system allows ease of changing reagents and uses min. organic solvent vols., as it avoids continuous pumping of reagents. Barbiturates (phenobarbital, amobarbital, pentobarbital and secobarbital) and serotonin re-uptake inhibitors (SRIs) - venlafaxine, paroxetine, sertraline and nortsertraline - were extracted as model acidic and basic compds. from urine into a 1:4 (v:v) mixture of 1-octanol and Bu chloride and back extracted into 0.45M NaOH and 0.18M H3PO4, resp. Absolute recoveries of the analytes were: venlafaxine 29; paroxetine 30; nortsertraline 25; sertraline 20; phenobarbital 11; amobarbital 24; pentobarbital 23; and secobarbital 27. The sample throughput, including extraction and back extraction, was 20 samples h⁻¹.
 ACCESSION NUMBER: 1996:752602 CAPLUS
 DOCUMENT NUMBER: 126:85763
 TITLE: Sequential-injection extraction for sample preparation
 AUTHOR(S): Peterson, Kristina L.; Logan, Barry K.; Christian, Gary D.; Ruzicka, Jaromir
 CORPORATE SOURCE: University of Washington, Department of Chemistry, Box 351700, Seattle, WA, 98195-1700, USA
 SOURCE: Analytica Chimica Acta (1997), 337(1), 99-106
 CODEN: ACACAM; ISSN: 0003-2670
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8P
 RL: ANT (Analyte); PUR (Purification or recovery); ANST (Analytical study); PREP (Preparation)
 (sequential-injection extraction for sample preparation)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 53 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




L4 ANSWER 54 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB A simple method for the measurement of sertraline and nortsertraline in plasma or serum suitable for use in single-dose pharmacokinetic studies has been developed. Internal standard solution, aqueous fenethazine (10 mg/L) (20 µL), and Tris buffer (2 mol/L, pH 10.6) (100 µL) were added to plasma (200 µL). Sertraline, nortsertraline and the internal standard were extracted into Me tert-Bu ether (200 µL) by mixing (30 s) and centrifugation (11,000 r.p.m., 4 min). A portion (100 µL) of the extract was injected onto a Spherisorb S5SCX HPLC column (150 + 4.6 mm i.d.) which was eluted with methanol:water (19 + 1) containing ammonium perchlorate (40 mmol/L), final pH 7.0. Detection was by UV monitoring (215 nm). The concentration of each analyte in each sample was calculated from the calibration graph (peak-height ratio of analyte to that of the internal standard against analyte concentration) obtained after anal. of plasma samples containing known amts. of sertraline and nortsertraline. The limit of accurate measurement of the assay was 10 µg/L sertraline and 20 µg/L nortsertraline.
 ACCESSION NUMBER: 1996:736563 CAPLUS
 DOCUMENT NUMBER: 126:69664
 TITLE: HPLC of sertraline and nortsertraline in plasma or serum
 AUTHOR(S): Patel, Jignasha; Spencer, E. P.; Flanagan, R. J.
 CORPORATE SOURCE: Poisons Unit., St. Thomas' Hosp. Trust, London, SE14 5ER, UK
 SOURCE: Biomedical Chromatography (1996), 10(6), 351-354
 CODEN: BICHE2; ISSN: 0269-3879
 PUBLISHER: Wiley
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 122873-19-2, Nortsertraline maleate
 RL: ANT (Analyte); BSU (Biological study, unclassified); MFM (Metabolic formation); ANST (Analytical study); BIOL (Biological study); FORM (Formation, nonpreparative)
 (sertraline and nortsertraline determination in plasma or serum by HPLC)
 RN 122873-19-2 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-, (2Z)-2-butenedioate (1:1) (9CI) (CA INDEX NAME)

CM 1
 CRN 87857-41-8
 CMF C16 H15 Cl2 N


Absolute stereochemistry.

L4 ANSWER 54 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



CM 2
 CRN 110-16-7
 CMF C4 H4 O4

Double bond geometry as shown.




L4 ANSWER 55 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB Venlafaxine (V) is a second-generation antidepressant approved for use in the United States in 1993. It is a derivative of phenethylamine and is structurally unrelated to first- and other second-generation antidepressants. Nevertheless, its mechanism of action is similar to other antidepressants; it inhibits the reuptake of presynaptic norepinephrine and serotonin. Its major routes of elimination involve O and N demethylation. O-Desmethylvenlafaxine (ODV) is biol. active. Therapeutic concns. of V and ODV are approx. 0.2 and 0.4 mg/L, resp. Three cases of drug intoxication involving V are presented. V and ODV were identified by gas chromatog.-nitrogen-phosphorus detection after alkaline extraction of the biol. specimen. On an HP-5 column, V and ODV elute after bupropion and fluoxetine, but prior to the first-generation antidepressants, sertraline, amoxapine, and trazodone. V and ODV were confirmed by full scan electron impact gas chromatog.-mass spectrometry. The heart-blood V and ODV concns. (mg/L) in the three cases were 6.6 and 31; 84 and 15, and 44 and 50, resp. In case 1, acetaminophen and diphenhydramine were found in the heart blood at 140 and 2.6 mg/L resp. In case 2, amitriptyline, nortriptyline, and chlordiazepoxide were found in the blood at 2.8, 0.5 and 3.3 mg/L, resp. In each case, the manner of death was suicide.

ACCESSION NUMBER: 1996:599888 CAPLUS
 DOCUMENT NUMBER: 125:240430
 TITLE: Distribution of venlafaxine in three postmortem cases
 AUTHOR(S): Levine, Barry; Jenkins, Amanda J.; Queen, Martin; Jufer, Rebecca; Smialek, John E.
 CORPORATE SOURCE: Office Chief Medical Examiner, Baltimore, MD, 21201, USA
 SOURCE: Journal of Analytical Toxicology (1996), 20(6), 502-505
 CODEN: JATOD3; ISSN: 0146-4760
 PUBLISHER: Preston Publications
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylsertraline
 RL: ANT (Analyte); ANST (Analytical study)
 (venlafaxine in three postmortem cases)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(SCI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 55 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




L4 ANSWER 56 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB N-demethylation of the selective serotonin reuptake inhibitor sertraline to desmethylsertraline yields a compound with 10- to 20-fold less potency at blocking serotonin (5-HT) reuptake as measured in vitro. In the present study desmethylsertraline (DMS) was examined in two in vivo models of reuptake inhibition-elevation of extracellular 5-HT in the corpus striatum as measured by microdialysis and inhibition of firing of serotonin-containing dorsal raphe neurons. Whereas sertraline (1, 3.2, and 10 mg/kg SC) produced a dose-dependent increase in extracellular 5-HT and a decrease in 5-HTAA in rat striatum, desmethylsertraline was without effect on either parameter. In similar fashion, desmethylsertraline had no effect on dorsal raphe cell firing at a dose (1,000 µg/kg IV) nearly 20-fold the ED50 for sertraline (52 µg/kg). Taken together, these data suggest that DMS does not contribute to the blockade of central 5-HT reuptake produced by sertraline in vivo and therefore would be expected to play a negligible role in its clin. activity.

ACCESSION NUMBER: 1996:239456 CAPLUS
 DOCUMENT NUMBER: 124:332755
 TITLE: Comparison of the effects of sertraline and its metabolite desmethylsertraline on blockade of central 5-HT reuptake in vivo
 AUTHOR(S): Sprouse, Jeffrey; Clarke, Thomas; Reynolds, Linda; Heym, James; Rollema, Hans
 CORPORATE SOURCE: Department Neuroscience, Pfizer Central Research, Groton, CT, 06340, USA
 SOURCE: Neuropsychopharmacology (1996), 14(4), 225-31
 CODEN: NEROEW; ISSN: 0893-133X
 PUBLISHER: Elsevier
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylsertraline
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative)
 (comparison of effects of sertraline and its metabolite desmethylsertraline on blockade of central 5-HT reuptake in vivo)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(SCI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 56 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



L4 ANSWER 57 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
AB Biotransformation of the triazolobenzodiazepine triazolam to its hydroxylated metabolites, α -hydroxy (OH)- and 4-OH-triazolam, was studied in vitro using microsomal prepn. of human liver. Mean values of V_{max} (10.3 nM/min/mg of protein) and K_m (304 μ M) for the 4-OH pathway exceeded values for the α -OH pathway (2.4 and 74, resp.). However, the mean V_{max}/K_m ratios for the two pathways were nearly identical, indicating that both contribute approx. equally to intrinsic clearance. Ketoconazole was a powerful inhibitor of triazolam biotransformation, having mean competitive K_i values of 0.006 and 0.023 μ M for the α -OH and 4-OH pathways. This is consistent with the role of P 450-3A isoforms in mediating triazolam biotransformation. The serotonin₂ antagonist antidepressant nefazodone inhibited the α -OH and 4-OH pathways (K_i = 0.6 and 1.7 μ M, resp.), but with considerably less activity than ketoconazole. Among six selective serotonin reuptake-inhibitor antidepressants, norfluoxetine was the most potent inhibitor (K_i = 2.7 and 8.0 μ M) and fluoxetine itself was the weakest (K_i = 7.0 and 44.3 μ M). In a double-blind clin. pharmacokinetic-pharmacodynamic study, administration of triazolam (0.125 mg) preceded by ketoconazole, compared to triazolam preceded by placebo, produced a

nearly 9-fold reduction in apparent oral clearance of triazolam (41 vs. 337 mL/min) and a 4-fold prolongation of half-life (13.5 vs. 3.4 h). Pharmacodynamic testing indicated enhancement of electroencephalog. beta activity, and enhanced decrements in digit-symbol substitution test performance, attributable to coadministration of ketoconazole. Plasma ketoconazole concns. measured in the clin. study ranged from 0.02 μ g/mL (projected min.) to 4.95 μ g/mL (maximum). An in vitro-in vivo scaling model, using these plasma ketoconazole concns. together with liver partition ratios

and the in vitro K_i values, predicted with liver partition ratios and the in vitro K_i values, predicted a decrement of triazolam clearance due to ketoconazole coadministration that was consistent with the 88% decrement in clearance activity observed in vivo.

ACCESSION NUMBER: 1996:112719 CAPLUS
DOCUMENT NUMBER: 124:219303
TITLE: Triazolam biotransformation by human liver microsomes in vitro: effects of metabolic inhibitors and

clinical conformation of a predicted interaction with ketoconazole
AUTHOR(S): von Moltke, Lisa L.; Greenblatt, David J.; Harmatz, Jerold S.; Duan, Su Xiang; Harrel, Lisa M.; Cotreau-Bibbo, Monette M.; Pritchard, Gary A.;

Wright, C. Eugene; Shader, Richard I.
CORPORATE SOURCE: Tufts University School of Medicine and The Division of Clinical Pharmacology, New England Medical Center, Boston, MA, USA


SOURCE: Journal of Pharmacology and Experimental Therapeutics (1996), 276(2), 370-9
CODEN: JPETAB; ISSN: 0022-3565

PUBLISHER: Williams & Wilkins
DOCUMENT TYPE: Journal
LANGUAGE: English

IT 87857-41-8, Desmethylertralin
RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)

L4 ANSWER 57 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
(metab. of triazolam and drug interactions)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 58 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
AB Desmethylertralin, a metabolite of the antidepressant drug sertraline, was compared with sertraline for its ability to produce effects characteristic of inhibitors of the serotonin transporter in vivo. Desmethylertralin antagonized brain serotonin depletion by p-chloroamphetamine, a depletion dependent upon the serotonin transporter, being less potent than sertraline in rats but almost as potent as sertraline in mice. Desmethylertralin was a weak antagonist of 6-hydroxydopamine-induced depletion of heart norepinephrine in mice; sertraline had no effect at the doses studied. Desmethylertralin decreased brain concns. of 5-hydroxyindoleacetic acid (5HIAA) in rats as did sertraline, the duration of the effect after both drugs being at

least 24 h but less than 48 h. After sertraline injection, desmethylertralin was present in rat brain at higher concns. than the parent drug at 8 h

and thereafter. In rats, repeated injections of sertraline, at doses previously shown to diminish β -adrenergic receptor-mediated responses, led to marked accumulation of desmethylertralin in brain and to inhibition of the catecholamine transporters. In mice, brain concns. of desmethylertralin were higher than those of parent drug within 7 h after sertraline injection and probably contributed importantly to the antagonism of p-chloroamphetamine effects. These data show that desmethylertralin is less potent than sertraline as a serotonin uptake inhibitor in vivo, as the in vitro data would have predicted, but that desmethylertralin may nonetheless contribute to the prolonged

inhibition of the serotonin transporter after sertraline administration, perhaps more in mice than in rats.

ACCESSION NUMBER: 1995:387035 CAPLUS
DOCUMENT NUMBER: 122:178161
TITLE: Comparison of desmethylertralin with sertraline as a

monoamine uptake inhibitor in vivo
AUTHOR(S): Fuller, Ray W.; Hemrick-Luecke, Susan K.; Littlefield, Ethel S.; Audia, James E.

CORPORATE SOURCE: Lilly Research Laboratories, Eli Lilly and Company, Indianapolis, IN, USA

SOURCE: Progress in Neuro-Psychopharmacology & Biological Psychiatry (1995), 19(1), 135-49
CODEN: PNPPD7; ISSN: 0278-5846

DOCUMENT TYPE: Journal
LANGUAGE: English


IT 87857-41-8, Desmethylertralin
RL: BAC (Biological activity or effector, except adverse); BPR (Biological process); BSU (Biological study, unclassified); MFM (Metabolic formation);

BIOL (Biological study); FORM (Formation, nonpreparative); PROC (Process) (comparison of desmethylertralin with sertraline as a monoamine uptake inhibitor in brain in vivo)

RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 58 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




L4 ANSWER 59 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB The biotransformation of the triazolobenzodiazepine alprazolam (ALP) to its hydroxylated metabolites (4-OH-ALP and α -OH-ALP) was evaluated in human, monkey, rat, and mouse liver microsomes. In all species 4-OH-ALP was the principal metabolite, accounting for 84% of clearance in human microsomes compared with 16% for α -OH-ALP. Among the serotonin-specific reuptake inhibitors fluoxetine (FLU) and sertraline (SERT), and their resp. demethylated metabolites norfluoxetine (NOR) and desmethylsertraline (DES), NOR was the most potent inhibitor (mean K_i for 4-OH-ALP formation in humans: 11 μ M), FLU the weakest (K_i = 83 μ M), with SERT and DES falling in between (K_i = 24 and 20 μ M). The in vitro data predict 29% inhibition of ALP clearance at mean FLU and NOR plasma concns. of 77 ng mL⁻¹ and 72 ng mL⁻¹, resp., after correction for liver:water partition ratios in the range of 12-14. The observed mean degree of inhibition in a previous in vivo study was 21%. Ketoconazole was a potent inhibitor of ALP metabolism in vitro (K_i = 0.046 μ M), suggesting that ALP hydroxylation is mediated by the cytochrome P 450-3A subfamily. Quinidine was a weak inhibitor (K_i = 626 μ M).

ACCESSION NUMBER: 1994:620997 CAPLUS
 DOCUMENT NUMBER: 121:220997
 TITLE: Inhibitors of alprazolam metabolism in vitro: Effect of serotonin-reuptake-inhibitor antidepressants, ketoconazole and quinidine
 AUTHOR(S): Moltke, Lisa L. von; Greenblatt, David J.; Cotreau-Bibbo, Monette M.; Harmatz, Jerold S.; Shader, Richard I.
 CORPORATE SOURCE: New England Medical Center Hospital, Tufts University School Medicine, Boston, MA, 02111, USA
 SOURCE: British Journal of Clinical Pharmacology (1994), 38(1), 23-31
 CODEN: BCPHEM; ISSN: 0306-5251
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylsertraline
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study) (alprazolam metabolism inhibition by desmethylsertraline)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 59 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




L4 ANSWER 60 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB Using radioligand binding assays and post-mortem normal human brain tissue, the authors obtained equilibrium dissociation consts. (K_{ds}) for 17 antidepressants and two of their metabolites at histamine H1, muscarinic, α_1 -adrenergic, α_2 -adrenergic, dopamine D2, serotonin 5-HT1A, and serotonin 5-HT2 receptors. Several newer antidepressants were compared with older drugs. In addition, the authors studied some antimuscarinic, antiparkinson, antihistamine, and neuroleptic compds. at some of these receptors. For the antidepressants, classical tricyclic antidepressants were the most potent drugs at five of the seven receptors (all but α_2 -adrenergic and 5-HT1A receptors). The chlorophenylpiperazine derivative antidepressants (etoperidone, nefazodone, trazodone) were the most potent antidepressants at α_2 -adrenergic and 5-HT1A receptors. Of ten antihistamines tested, none was more potent than doxepin at histamine H1 receptors. At muscarinic receptors antidepressants and antihistamines had a range of potencies, which were mostly weaker than those for antimuscarinics. From the in vitro data, the authors expect adinazolam, bupropion, fluoxetine, sertraline, tomoxetine, and venlafaxine not to block any of these five receptors in vivo. An antidepressant's potency for blocking a specific receptor is predictive of certain side effects and drug-drug interactions. These studies can provide guidelines for the clinician in the choice of antidepressant.

ACCESSION NUMBER: 1994:449972 CAPLUS
 DOCUMENT NUMBER: 121:49972
 TITLE: Binding of antidepressants to human brain receptors: focus on newer generation compounds
 AUTHOR(S): Cusack, Bernadette; Nelson, Albert; Richelson, Elliott
 CORPORATE SOURCE: Dep. Res., Mayo Clin. Jacksonville, Jacksonville, FL, 32224, USA
 SOURCE: Psychopharmacology (Berlin, Germany) (1994), 114(4), 559-65
 CODEN: PSCHDL; ISSN: 0033-3150
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylsertraline
 RL: PROC (Process) (binding of, to human brain receptors)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.


L4 ANSWER 60 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



L4 ANSWER 61 OF 79 CAPLUS COPYRIGHT 2004 ACS ON STN
 AB The authors report methods for the anal. of sertraline and desmethylsertraline in postmortem biol. fluids. The extraction method is based on a widely used procedure employing Bu chloride, and instrumental anal. is performed using GC/MS and HPLC with photodiode array detection. The authors report retention index, mass spectral, and UV-vis properties of the drug and its metabolite. Samples from three sertraline-related deaths were analyzed and revealed concns. up to 10 times greater than the normal therapeutic levels, although two of the deaths were obviously the result of other causes. The authors also noted in one case that the drug concns. in central and peripheral blood were very similar, suggesting that postmortem distribution may be uniform.

ACCESSION NUMBER: 1994:426134 CAPLUS
 DOCUMENT NUMBER: 121:26134
 TITLE: Analysis of sertraline (Zoloft) and its major metabolite in postmortem specimens by gas and liquid chromatography
 AUTHOR(S): Logan, B. K.; Friel, P. N.; Case, G. A.
 CORPORATE SOURCE: Dep. Lab. Med., Univ. Washington, Seattle, WA, 98134, USA
 SOURCE: Journal of Analytical Toxicology (1994), 18(3), 139-42
 CODEN: JATOD3; ISSN: 0146-4760
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylsertraline
 RL: ANT (Analyte); ANST (Analytical study)
 (determination of, as sertraline metabolite, in postmortem human blood by gas and liquid chromatog.)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 62 OF 79 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)

L4 ANSWER 62 OF 79 CAPLUS COPYRIGHT 2004 ACS ON STN
 AB The determination of the new antidepressant drug sertraline and its main metabolite, desmethylsertraline, in human serum is described. A new solid-phase extraction method employing the dual functionality Clean Screen® cartridge is presented followed by reversed-phase liquid chromatog. (LC) anal. The sample preparation yielded extremely clean exts. and absolute recoveries in excess of 90% for both drugs from human serum. The response of the LC system was linear over the concentration range 0.01-2.5mg/L for both sertraline and desmethylsertraline with a limit of detection of 0.01mg/L. A gas chromatog. mass spectrometric (GC-MS) system is also described should confirmation of the drugs be necessary.

ACCESSION NUMBER: 1994:289312 CAPLUS
 DOCUMENT NUMBER: 120:289312
 TITLE: Determination of sertraline and desmethylsertraline in human serum using copolymeric bonded-phase extraction, liquid chromatography and gas chromatography-mass spectrometry
 AUTHOR(S): Rogowsky, Dawn; Marr, Margaret; Long, Gerald; Moore, Christine
 CORPORATE SOURCE: United Chemical Technologies, 2731 Bartram Road, Bristol, PA, 19007, USA
 SOURCE: Journal of Chromatography, B: Biomedical Sciences and Applications (1994), 655(1), 138-41
 CODEN: JCBREP; ISSN: 1387-2273
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylsertraline
 RL: ANT (Analyte); ANST (Analytical study)
 (determination of, in human blood by reversed-phase liquid chromatog. and gas chromatog.-mass spectrometry)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 63 OF 79 CAPLUS COPYRIGHT 2004 ACS ON STN
 AB Biotransformation of the tricyclic antidepressant desipramine (DMI) to its metabolite 2-hydroxy-desipramine (2-OH-DMI) was studied in vitro using microsomal preps. from human, monkey, mouse and rat liver. In all species 2-OH-DMI was the principal identified metabolite. Mean (±S.E.) reaction parameters in six human liver samples were: Vmax, 0.11 ± 0.02 nmol/mL/min/mg protein; Km, 16.1 ± 4.2 µM. Quinidine was a highly potent inhibitor of 2-OH-DMI formation (mean K_i = 0.053 µM), consistent with the presumed role of Cytochrome P 450-2D6 in mediating this reaction. Ketoconazole was a much less potent inhibitor (mean K_i = 10.3 µM). Two serotonin-specific reuptake inhibitor (SSRI) antidepressants, and their resp. metabolites, were evaluated as potential inhibitors of 2-OH-DMI formation. Fluoxetine (FLU) and norfluoxetine (NOR) were the most potent inhibitors (mean K_i values: 3.0 and 3.5 µM, resp.). Sertraline (SERT) and its metabolite desmethylsertraline (DES) also inhibited the reaction (mean K_i: 22.7 and 16.0 µM), but were significantly less potent than FLU or NOR. Values of K_i and K_m measured in vitro were used to generate a theor. prediction of the degree of clearance inhibition in vivo at any given concentration of substrate and inhibitor. The model was applied to a clin. study in which DMI clearance in humans was impaired by coadministration of FLU (yielding FLU and NOR in plasma) or by SERT (yielding SERT and DES in plasma). Use of plasma SSRI concns. in the predictive model underestimated the actual impairment of DMI clearance. However, in vitro liver:water partition ratios for the SSRIs were in the range of 12 to 14, suggesting that liver concns. will considerably exceed those in plasma. Use of projected liver SSRI concns. (plasma level + partition ratio) in the model yielded highly accurate predictions. Predicted vs. observed impairments of DMI clearance were: 81.7 vs. 79.2% due to FLU/NOR; 17.5 vs. 16.7% due to SERT/DES. Thus, in vitro drug metabolism models can provide qual. and quant. data on drug interactions that are of direct clin. relevance.

ACCESSION NUMBER: 1994:260419 CAPLUS
 DOCUMENT NUMBER: 120:260419
 TITLE: Inhibition of desipramine hydroxylation in vitro by serotonin-reuptake-inhibitor antidepressants, and by quinidine and ketoconazole: a model system to predict drug interactions in vivo
 AUTHOR(S): Von Moltke, Lisa L.; Greenblatt, David J.; Cotreau-Bibbo, Monette M.; Duan, Su Xiang; Harmatz, Jerold S.; Shader, Richard I.
 CORPORATE SOURCE: Sch. Med., Tufts Univ., Boston, MA, USA
 SOURCE: Journal of Pharmacology and Experimental Therapeutics (1994), 268(3), 1278-83
 CODEN: JPETAB; ISSN: 0022-3565
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylsertraline
 RL: ANST (Analytical study)
 (desipramine hydroxylation in liver microsome model inhibition by, drug interaction prediction in relation to)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

L4 ANSWER 63 OF 79 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)

Absolute stereochemistry.




L4 ANSWER 64 OF 79 CAPLUS COPYRIGHT 2004 ACS ON STN

AB Paroxetine is a selective and potent inhibitor of 5-hydroxytryptamine (5-HT) uptake into serotonergic neurons. [3H]paroxetine binding to rat frontal cortex was of high affinity with a high percentage of sp. binding. The binding data of both competition and saturation studies fitted a single site binding model. [3H]paroxetine binding was potentially inhibited by the selective 5-HT uptake inhibitors. In addition, a very good correlation was demonstrated between the ability of twenty-three compds. to inhibit [3H]paroxetine binding to rat frontal cortical membranes and [3H]5-HT uptake into rat frontal cortical synaptosomes. These data support the view that [3H]paroxetine binds to a single site which corresponds to the 5-HT uptake site. Using this ligand, the effects of repeated administration of antidepressant drugs with a wide range of pharmacol. actions and electroconvulsive shock on 5-HT reuptake sites were examined [3H]paroxetine binding parameters (Kd and Bmax) were unaltered by all treatments. It would, therefore, appear that antidepressant therapy does not produce adaptive changes in 5-HT uptake sites.

ACCESSION NUMBER: 1993:595588 CAPLUS
DOCUMENT NUMBER: 119:195588
TITLE: [3H]Paroxetine binding in rat frontal cortex strongly correlates with [3H]5-HT uptake: effect of administration of various antidepressant treatments
AUTHOR(S): Cheetham, S. C.; Viggers, J. A.; Slater, N. A.; Heal, D. J.; Buckett, W. R.
CORPORATE SOURCE: Boots Pharm. Res. Dep., Nottingham, NG2 3AA, UK
SOURCE: Neuropharmacology (1993), 32(8), 737-43
CODEN: NEPHW; ISSN: 0028-3908
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 122873-19-2, Desmethylertraline maleate
RL: BIOL (Biological study)
(hydroxytryptamine uptake inhibition by, paroxetine binding inhibition in relation to, in brain frontal cortex)
RN 122873-19-2 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-, (2Z)-2-butenedioate (1:1) (9CI) (CA INDEX NAME)
CM 1
CRN 87857-41-8
CMF C16 H15 Cl2 N


Absolute stereochemistry.

L4 ANSWER 64 OF 79 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



CM 2
CRN 110-16-7
CMF C4 H4 O4

Double bond geometry as shown.




L4 ANSWER 65 OF 79 CAPLUS COPYRIGHT 2004 ACS ON STN

AB Based on mol. cloning studies, five different muscarinic receptor subtypes exist: m1, m2, m3, m4, and m5. The authors determined the affinity and selectivity of binding for sixteen antidepressants, two of their metabolites, and three antihistaminics (H1) at these subtypes. Using Chinese hamster ovary cells (CHO-K1) transfected with genes for the human muscarinic receptor subtypes, the authors obtained equilibrium dissociation constants. (Kds) from competitive radioligand binding studies with [3H]-quinuclidinyl benzilate ([3H]QNB) and membrane preps. of these cells. QNB was the most potent compound studied (Kd 30-80 pM). Mequitazine (Kd 6-14 nM) and amitriptyline (Kd 7-16 nM) exhibited the highest affinity among the antihistaminics and antidepressants, resp. Among the antidepressants examined were the serotonin-selective drugs sertraline and fluoxetine, both of which displayed Kd values >1 μM. The remaining antidepressants were moderate to weak antagonists with some eliciting no radioligand competition at high concns. The compds. studied showed no significant selectivity among the five cloned subtypes.

ACCESSION NUMBER: 1993:531407 CAPLUS
DOCUMENT NUMBER: 119:131407
TITLE: Antagonism of the five cloned human muscarinic cholinergic receptors expressed in CHO-K1 cells by antidepressants and antihistaminics
AUTHOR(S): Stanton, Tiffany; Bolden-Watson, Carolyn; Cusack, Bernadette; Richelson, Elliott
CORPORATE SOURCE: Dep. Psychiatry, Mayo Found. and Mayo Clin., Jacksonville, FL, 32224, USA
SOURCE: Biochemical Pharmacology (1993), 45(11), 2352-4
CODEN: BCPCA6; ISSN: 0006-2952
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylertraline
RL: PROC (Process)
(binding of, to human muscarinic receptor subtypes, expressed in Chinese hamster ovary cells)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-, (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 66 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB We determined the uptake blockade produced by eight new antidepressant drugs (etoperidone, femoxetine, lofepramine, nefazodone, paroxetine, sertraline, tomoxetine, and venlafaxine), two metabolites of newer antidepressants, and carbamazepine. Inhibitor consts. (Kis) for uptake blockade were obtained from competitive uptake studies with [3H]norepinephrine, [3H]5-hydroxytryptamine, and [3H]dopamine in rat brain synaptosomes prepared from hippocampus, frontal cortex, and striatum, resp. Among the newer compds., tomoxetine (Ki = 0.7 nM) and lofepramine (Ki = 1.9 nM) were potent and selective [3H]norepinephrine uptake blockers; paroxetine (Ki = 0.73 nM), sertraline (Ki = 3.4 nM), and femoxetine (Ki = 22 nM) potently and selectively inhibited [3H]5-hydroxytryptamine uptake. Although none of the drugs was potent for [3H]dopamine uptake blockade, sertraline was the most potent (Ki = 260 nM). These data are useful in predicting adverse effects and drug-drug interactions of antidepressants.

ACCESSION NUMBER: 1993:183305 CAPLUS
 DOCUMENT NUMBER: 118:183305
 TITLE: Blockade by newly-developed antidepressants of biogenic amine uptake into rat brain synaptosomes
 AUTHOR(S): Bolden-Watson, C.; Richelson, E.
 CORPORATE SOURCE: Mayo Clin., Jacksonville, FL, 32224, USA
 SOURCE: Life Sciences (1993), 52(12), 1023-9
 CODEN: LIFSAK; ISSN: 0024-3205

DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8, Desmethylertraline
 RL: BIOL (Biological study)
 (biogenic amine uptake inhibition by, in brain synaptosomes, antidepressant activity in relation to)

RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 67 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB Some agents that increase serotonergic transmission in the brain show anorectic activity in doses that do not interfere with the behavior of rats and other animal species. These agents reduce food intake by a mechanism that clearly differs from that involved in the anorectic activity of d-amphetamine. d-Fenfluramine, fluoxetine, and sertraline are three drugs that have already been tested and are used in man. These compds. accumulate in the brain and are metabolized through N-dealkylation. They affect the uptake and release of serotonin at different concns., with mechanisms that do not completely overlap. There is pharmacol. evidence that d-fenfluramine and sertraline exert their anorectic activity by enhancing the stimulation of 5-HT1nonA receptors whereas fluoxetine seems to affect at anorectic doses both serotonergic and dopaminergic systems. The role of serotonin in controlling food intake will be discussed, and the effects of agents that reduce serotonergic transmission will also be considered.

ACCESSION NUMBER: 1992:228034 CAPLUS
 DOCUMENT NUMBER: 116:228034
 TITLE: Progress report on the anorexia induced by drugs believed to mimic some of the effects of serotonin on the central nervous system
 AUTHQR(S): Garattini, Silvio; Bizzzi, Adalgisa; Codegani, Anna Maria; Caccia, Silvio; Mennini, Tiziana
 CORPORATE SOURCE: Ist. Ric. Farmacol. "Mario Negri", Milan, 20157, Italy
 SOURCE: American Journal of Clinical Nutrition (1992), 55(1, Suppl.), 160S-166S
 CODEN: AJCNAC; ISSN: 0002-9165

DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study) (sertraline metabolite, anorectic activity of, brain serotonergic neurotransmission modulation in)

RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 69 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
AB A quant. structure-activity relationship (QSAR) was analyzed for
monoamine
uptake inhibition of cis- and trans-1-alkylamino-4-aryltetralins using
the
Free-Wilson method. The calculated individual substituent contributions
were

strongly correlated with the uptake inhibition activity of dopamine,
5-hydroxytryptamine and norepinephrine in the corpus striatum and
hypothalamus of rats, resp. The effect of the substituents and
stereochem. factors on the monoamine uptake inhibition of the analyzed
series of tetralins was discussed.

ACCESSION NUMBER: 1991:526385 CAPLUS
DOCUMENT NUMBER: 115:126385
TITLE: Structure-activity relationship studies of CNS
agents.


AUTHOR(S): Part II. De novo analysis of the monoamine uptake
CORPORATE SOURCE: Mokrosz, Jerzy L.
Inst. Pharmacol., Pol. Acad. Sci., Krakow, 31-343,
Pol.

SOURCE: Polish Journal of Pharmacology and Pharmacy (1990),
42 (5), 501-7
CODEN: PJPPAA; ISSN: 0301-0244

DOCUMENT TYPE: Journal
LANGUAGE: English
IT 91797-58-9


RL: BIOL (Biological study)
(monoamines uptake by brain inhibition by, structure in relation to)
RN 91797-58-9 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, cis- (9CI)
(CA INDEX NAME)

Relative stereochemistry.




L4 ANSWER 69 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
IT 87857-41-8
RL: BIOL (Biological study)
(pharmaceutical containing, as inflammation and psychosis inhibitor
and
immunosuppressant)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 69 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
GI



AB The title compds. I [R1,R2 = H, Cl-3 alkyl; Z = (un)substituted Ph; W =
H,

F, Br, Cl, CF3, alkoxy] are drugs for treatment of schizophrenia,
psychosis, mania, etc., and are also useful as immunosuppressants and
inflammation inhibitors. (+)-trans-I (R1 = W = H; R2 = Me; Z =
4-FC6H4) inhibited the in vitro binding of (+)-3-(3-hydroxyphenyl)-N-(1-
propyl)piperidine-3H to the α receptors of the rat brain.

ACCESSION NUMBER: 1991:221410 CAPLUS
DOCUMENT NUMBER: 114:221410
TITLE: 4-Phenyl-1,2,3,4-tetrahydro-1-naphthalenamine
derivatives in the treatment of psychosis,
inflammation, stroke and as immunosuppressants
Koe, Kenneth B.
Pfizer Inc., USA
Eur. Pat. Appl., 8 pp.
CODEN: EPXKDW

DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 386997	A2	19900912	EP 1990-302365	19900306
EP 386997	A3	19920429		
EP 386997	B1	19970319		
US 4981870	A	19910101	US 1989-320014	19890307
IL 93576	A1	19951208	IL 1990-93576	19900228
CA 2011428	AA	19900907	CA 1990-2011428	19900305
CA 2011428	C	19941108		
AU 9050796	A1	19900920	AU 1990-50796	19900306
AU 610036	B2	19910509		
JP 02300121	A2	19901212	JP 1990-54802	19900306
JP 07014870	B4	19950222		
HU 59000	A2	19920428	HU 1990-1325	19900306
AT 150299	E	19970415	AT 1990-302365	19900306
ZA 9001743	A	19911030	ZA 1990-1743	19900307
US 5061728	A	19911029	US 1990-622308	19901205
JP 07173056	A2	19950711	JP 1994-172929	19940725
PRIORITY APPLN. INFO.:			US 1989-320014	A 19890307


OTHER SOURCE(S): MARPAT 114:221410

L4 ANSWER 70 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
AB The effects of repeated treatment with the serotonin uptake blocker
sertraline on cocaine-induced locomotion in female C57Bl/6BYJ mice were
examined in three paradigms. First, when animals were treated for 2 wk
with
a daily injection of 8 mg/kg i.p. of sertraline (or placebo) and
challenged with cocaine (25 mg/kg i.p.) 1 h after the final sertraline
injection, their cocaine-induced locomotion was the same as that of
placebo-pretreated controls. Second, animals were treated for 2 wk with
cocaine (25 mg/kg i.p. once a day) (or saline) and then for 2 wk with
sertraline (8 mg/kg i.p. once a day) (or placebo). Locomotion induced by
cocaine (25 mg/kg i.p.) administered 1 h after the final sertraline
(placebo) injection was higher in cocaine- than saline-pretreated mice
(sensitization), but there was no difference between sertraline- and
placebo-pretreated animals. Third, daily treatment with sertraline (8
mg/kg i.p.) did not change the locomotor stimulatory effect of cocaine
(25
mg/kg i.p.) administered after a 3-wk continuous infusion of cocaine (22
mg/kg/day s.c.) by osmotic minipumps or after three, four, or seven
injections of cocaine (15 or 25 mg/kg i.p.). After cocaine
administration
(25 mg/kg i.p.), animals pretreated repeatedly with sertraline (8 mg/kg
i.p.
once a day for 2 wk) had the same plasma or brain levels of cocaine as
those pretreated with placebo; there was no difference between cocaine-
and saline-treated mice in brain levels of sertraline or
desmethylsertraline.

ACCESSION NUMBER: 1991:157119 CAPLUS
DOCUMENT NUMBER: 114:157119
TITLE: Sertraline and cocaine-induced locomotion in mice.
II. Chronic studies
Reith, Marten E. A.; Fischette, Christine T.
Cent. Neurochem., N.S. Kline Inst. Psychiatr. Res.,
New York, NY, 10035, USA
SOURCE: Psychopharmacology (Berlin, Germany) (1991), 103(3),
306-13
CODEN: PSCHDL; ISSN: 0033-3158
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: BIOL (Biological study)
(as sertraline metabolite, cocaine effect on)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.

L4 ANSWER 70 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)




L4 ANSWER 71 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN

AB The present study assessed the behavioral and pharmacokinetic interaction between the serotonin uptake blocker sertraline and cocaine in C57BL/6ByJ mice. Pretreatment with sertraline (1-32 mg/kg i.p.) did not affect the total amount of spontaneous locomotor activity during 50 min following administration of cocaine (15-40 mg/kg i.p.). At doses of sertraline (16 and 32 mg/kg) much higher than those found to inhibit *ex vivo* neuronal uptake of serotonin by 50% (1-2 mg/kg), the peak of cocaine-induced locomotor activity was shifted towards a later time. A similar effect


was seen after pretreatment with serotonin uptake blockers other than sertraline, and also after desipramine. Sertraline (16 and 32 mg/kg), given 60 min prior to cocaine, did not affect levels of cocaine in brain and plasma, and cocaine administration did not alter the brain level of sertraline. Although female mice were more responsive to cocaine than male mice, they were not different in their response to sertraline.

ACCESSION NUMBER: 1991:157118 CAPLUS
DOCUMENT NUMBER: 114:157118
TITLE: Sertraline and cocaine-induced locomotion in mice.
I. Acute studies
AUTHOR(S): Reith, Maarten E. A.; Wiener, Harvey L.; Fischette, Christine T.
CORPORATE SOURCE: Cent. Neurochem., N.S. Kline Inst. Psychiatr. Res., New York, NY, 10035, USA
SOURCE: Psychopharmacology (Berlin, Germany) (1991), 103(3), 297-305
CODEN: PSCHDL; ISSN: 0033-3158
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: BIOL (Biological study)
(as sertraline metabolite, cocaine effect on)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 72 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
GI




AB A rapid isocratic reversed-phase HPLC method for simultaneous determination of sertraline (I) and its metabolite desmethylsertraline (II) in mouse cerebral cortex is described. The method is based on a Versapak C18 10- μ m-column, MeCN-0.25M KH₂PO₄ (pH 2.7) buffer (30:70), the flow rate of 2 mL/min, and tetracaine as an internal standard. The retention time was 9.5 and 12 min for II and I, resp. Standard curves for I and II were linear and had the following regression coeffs.: for I: slope = 0.105, y-axis intercept = -0.8, r^2 = 0.998; for II: slope = 0.11, y-axis intercept = 1.37, r^2 = 0.989. After one extraction with EtOH, 89% of the I and 87% of the II was recovered. The coefficient of variations for I and II was 4.8 and 4.9%, resp. After i.p. administration of 32 mg/kg I, the drug was detectable as early as 5 min after injection. The metabolite was first observed 15 min after administration of I. The detection limit was approx. 20 and 10 pmol/mg wet weight for I and II, resp.


ACCESSION NUMBER: 1990:508607 CAPLUS
DOCUMENT NUMBER: 113:108607
TITLE: Separation and determination of sertraline and its metabolite, desmethylsertraline, in mouse cerebral cortex by reversed-phase high-performance liquid chromatography
AUTHOR(S): Wiener, Harvey L.; Kramer, H. Kenneth; Reith, Maarten E. A.
CORPORATE SOURCE: Coll. Pharm. Allied Health Prof., St. John's Univ., Jamaica, NY, 11439, USA
SOURCE: Journal of Chromatography (1990), 527(2), 467-72
CODEN: JOCRAM; ISSN: 0021-9673
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8, Desmethylsertraline
RL: ANT (Analyte); ANST (Analytical study)
(determination of, as sertraline metabolite, in cerebral cortex, by reversed-phase HPLC)
RN 87857-41-8 CAPLUS

L4 ANSWER 72 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.



L4 ANSWER 73 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
GI



AB Sertraline (I) is a potent and selective inhibitor of neuronal serotonin uptake and is currently under development for the treatment of depression and of obesity. The drug was >97% bound to plasma proteins, yet extensively distributed into tissues. The whole brain concentration of sertraline in the rat was >40-fold higher than that in plasma, and the volume of distribution was about 25 L/kg in the rat and dog. Sertraline

was extensively metabolized by the rat and dog prior to excretion. The metabolic clearance of sertraline was >35 mL of blood/min/kg in each species, and 1st-pass metabolism occurred with oral administration.

Initial metabolic steps included N-demethylation, N-hydroxylation, oxidative deamination, and glucuronidation of sertraline carbamic acid, which in solution was in equilibrium with sertraline and CO₂. The N-demethyl metabolite, which was 10-fold less potent as an inhibitor of serotonin uptake, was formed in both species. Plasma area under the concentration-time curve for demethylsertraline was 66-270% of that for sertraline, and was dependent on the species examined and route of drug administration. Sertraline and demethylsertraline underwent oxidative deamination to the corresponding ketone, which was subsequently hydroxylated at the α-carbon, forming a diastereomeric metabolite pair. The glucuronides of sertraline carbamic acid, N-hydroxysertraline, and the α-hydroxy ketone diastereomers comprised 45% and 82% of the total radiolabel excreted in urine plus bile of duct-cannulated rats and dogs, resp. Bile was the major route of elimination in both species.


ACCESSION NUMBER: 1990:15883 CAPLUS
DOCUMENT NUMBER: 112:15883
TITLE: Metabolism and disposition of the 5-hydroxytryptamine uptake blocker sertraline in the rat and dog
AUTHOR(S): Tremaine, Larry M.; Welch, Willard M.; Ronfeld, Robert
CORPORATE SOURCE: A. Drug Metab. Dep., Pfizer, Inc., Groton, CT, USA
SOURCE: Drug Metabolism and Disposition (1989), 17(5), 542-50
CODEN: DMDSAI; ISSN: 0090-9556
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 87857-41-8P, Demethylsertraline
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and formation of, as sertraline metabolite)

L4 ANSWER 74 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
AB [N-methyl-¹¹C]Sertraline, a potential agent for the study of the serotonergic system in vivo with positron emission tomog., was prepared by N-methylation of the corresponding norcompound with [¹¹C]iodomethane, which was itself prepared from cyclotron-produced [¹¹C]carbon dioxide. Under the best conditions found [norsertraline free base (20 mM) in DMF (0.70 mL), 120° for 8 min] [N-methyl-¹¹C]sertraline can be prepared in 43% radiochem. yield from [¹¹C]iodomethane (decay-corrected), corresponding to 20% overall radiochem. yield from [¹¹C]carbon dioxide (decay-corrected), with high specific radioactivity. Preps. can be ready for i.v. injection 50 min from the end of radionuclide production

ACCESSION NUMBER: 1989:55324 CAPLUS
DOCUMENT NUMBER: 111:153324
TITLE: The radiosynthesis of [N-methyl-¹¹C]-sertraline
AUTHOR(S): Lasne, M. C.; Pike, V. W.; Turton, D. R.
CORPORATE SOURCE: MRC Cyclotron Unit, Hammersmith Hosp., London, W12 0HS, UK
SOURCE: Applied Radiation and Isotopes (1989), 40(2), 147-51
CODEN: ARISEF; ISSN: 0883-2889
DOCUMENT TYPE: Journal
LANGUAGE: English
IT 122873-19-2, Norsertraline maleate
RL: PROC (Process)
(conversion of, to norsertraline free base)
RN 122873-19-2 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-, (2Z)-2-butenedioate (1:1) (9CI) (CA INDEX NAME)

CM 1

CRN 87857-41-8
CMF C16 H15 Cl2 N




CM 2

CRN 110-16-7


L4 ANSWER 73 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)- (9CI) (CA INDEX NAME)

Absolute stereochemistry.




L4 ANSWER 74 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
CMF C4 H4 O4

Double bond geometry as shown.



IT 87857-41-8P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation of)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)- (9CI) (CA INDEX NAME)


Absolute stereochemistry.




L4 ANSWER 75 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB QSAR of monoamine uptake inhibition by cis- and trans-1-amino-4-aryltetralins was examined. Physicochem. parameters such as electronic (σ), hydrophobic (π), and molar refraction (MR) were significantly correlated with 5-HT [50-67-9] uptake inhibition in corpus striatum of rat brain for both series of compds. The norepinephrine (NE) [51-41-2] uptake inhibition in hypothalamus and dopamine (DA) [51-61-6] uptake inhibition in corpus striatum, however, yielded poor correlations with either set of compds. However, the selectivity ratios, $-\log [IC_{50} (DA)/IC_{50} (5-HT)]$ and $-\log [IC_{50} (NE)/IC_{50} (5-HT)]$ described statistically sound correlations with properly chosen physicochem. parameters for cis- and trans-1-amino-4-aryltetralins.

ACCESSION NUMBER: 1987:148963 CAPLUS
 DOCUMENT NUMBER: 106:148963
 TITLE: Quantitative structure-activity relationship of antidepressant agents. Derivatives of cis- and trans-1-amino-4-aryltetralin
 Singh, P.; Goyal, A.
 CORPORATE SOURCE: Dep. Chem., Birla Inst. Technol. Sci., Pilani, India
 SOURCE: Arzneimittelforschung (1987), 37(1), 51-4
 CODEN: ARZNAD; ISSN: 0004-4172
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 91797-58-9
 RL: BIOL (Biological study)
 (brain monoamine uptake inhibition by, QSAR of)
 RN 91797-58-9 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, cis- (9CI)
 (CA INDEX NAME)

Relative stereochemistry.



L4 ANSWER 76 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
 Absolute stereochemistry.




L4 ANSWER 76 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 AB The effects of 2 specific serotonin re-uptake inhibitors, sertraline [79617-96-2] and Organic 6582 [59905-71-4], on the behavior of the bilaterally olfactory bulbectomized rat in the open field apparatus were compared following their acute and chronic administration. Neither drug affected the increased ambulation scores of the bulbectomized rats following their acute administration, but both drugs significantly attenuated the hypermotility of the bulbectomized rats after chronic (12 d) administration. Only Organic 6582 reduced the increased rearing score in the open field are qual. similar to that found in other studies for both typical (e.g., amitriptyline) and atypical (e.g., mianserin) antidepressants. The major metabolite of sertraline, CP-53261 (desmethylsertraline) [87857-41-8] was inactive in this behavioral test. The effects of sertraline on the behavior of rats in the hole board apparatus was also assessed. Sertraline, after both its acute and chronic administration, increased the number of crossings of the bulbectomized rats, without affecting the sham-operated controls. Unlike the stressful open field apparatus, the less stressful hole board apparatus cannot be used to differentiate between the bulbectomized rats, their sham-operated controls and the chronic effect of antidepressants.

Organic 6582 increased the serotonin [50-67-9] content of the mid-brain of the bulbectomized rats and reduced that of 5-HIAA, without appreciably affecting the concns. of GABA [56-12-2], noradrenaline [51-41-2] and serotonin in the amygdaloid cortex or mid-brain. By contrast, sertraline increased the dopamine [51-61-6] concentration in the amygdaloid cortex of the sham-operated and bulbectomized rats and raised the serotonin content of the mid-brain without affecting that of 5-HIAA. Apparently the attenuation of the hypermotility in the open field apparatus might be correlated with a rise in the serotonin content in this region. The relation of these results to the antidepressant activity of sertraline and Organic 6582 and the use of olfactory bulbectomized rats as an animal model of depression are discussed.

ACCESSION NUMBER: 1986:199968 CAPLUS
 DOCUMENT NUMBER: 104:199968
 TITLE: Effect of two specific serotonin re-uptake inhibitors on the behavior of the olfactory bulbectomized rat in the 'open field' apparatus
 Earley, Bernadette; Leonard, B. E.
 CORPORATE SOURCE: Pharmacol. Dep., Univ. Coll. Galway, Galway, Ire.
 SOURCE: Biological Psychiatry--New Prospects (1985), 5(Clin. Pharmacol. Stud. Psychiatr. Disord.), 234-40
 CODEN: BPNPES; ISSN: 0266-2124
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8
 RL: BIOL (Biological study)
 (hyperkinesia in olfactory bulbectomized rat response to, antidepressant activity in relation to, as sertraline metabolite)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)- (9CI) (CA INDEX NAME)


L4 ANSWER 77 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN
 GI



AB The title compound enantiomers I (R1 and R2 = H or Me; R3 = H, Cl, or MeO; R4 = H, Cl, CF3, or MeO; R5 = H, Br, Cl, F, CF3, MeO, BuO, or PhO) mostly as the HCl salts were prepared from the appropriate benzophenone and 136 1-tetralone [52738-06-2] and evaluated in vitro for their ability to inhibit the uptake of dopamine and serotonin in corpus striatum and of epinephrine in hypothalamus of rats. The cis compds. are potent and selective inhibitors of serotonin uptake, whereas the trans compds. block uptake of dopamine and norepinephrine. Structure-activity relations are discussed.


ACCESSION NUMBER: 1984:622093 CAPLUS
 DOCUMENT NUMBER: 101:222093
 TITLE: Nontricyclic antidepressant agents derived from cis- and trans-1-amino-4-aryltetralins
 Welch, Willard M.; Kraska, Allen R.; Sarges, Koe, B. Kenneth
 CORPORATE SOURCE: Cent. Res. Div., Pfizer Inc., Groton, CT, 06340, USA
 SOURCE: Journal of Medicinal Chemistry (1984), 27(11), 1508-15
 CODEN: JMCMMR; ISSN: 0022-2623
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 101:222093
 IT 91797-58-9
 RL: BAC (Biological activity or effector, except adverse); BSU (Biological study, unclassified); BIOL (Biological study)
 (monoamine uptake by brain response to)
 RN 91797-58-9 CAPLUS
 CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, cis- (9CI)
 (CA INDEX NAME)

Relative stereochemistry.




IT 91797-57-8P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (preparation and monoamine uptake by brain response to)
 RN 91797-57-8 CAPLUS
 CN 1-Naphthalenamine, 4-[(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-,
 hydrochloride, cis- (9CI) (CA INDEX NAME)

Relative stereochemistry.





● HCl



AB The pharmacol. activity of sertraline (I) [79617-96-2] was evaluated in tests predictive of 5-HT [50-67-9] uptake inhibiting and antidepressant activity. I is a highly selective and potent inhibitor of synaptosomal 5-HT uptake in rat brain. N-Demethylsertraline [87857-41-8], the major metabolite of I, is also a selective 5-HT uptake blocker, indicating that this selective inhibition is probably maintained in vivo. I markedly reduced the immobility of mice in the Porsolt forced swimming test; in addition repeated dosing of I induced down regulation of norepinephrine coupled adenylate cyclase in rat limbic forebrain. Thus,

I should be clin. effective as an antidepressant.
 ACCESSION NUMBER: 1984:45134 CAPLUS
 DOCUMENT NUMBER: 100:45134
 TITLE: Sertraline: a new selective inhibitor of serotonin uptake
 AUTHOR(S): Koe, B. Kenneth; Weissman, Albert; Welch, Willard M.; Browne, Ronald G.
 CORPORATE SOURCE: Cent. Res. Div., Pfizer Inc., Groton, CT, 06340, USA
 SOURCE: Psychopharmacology Bulletin (1983), 19(4), 687-91
 CODEN: PSYBB9; ISSN: 0048-5764
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 IT 87857-41-8
 RL: BIOL (Biological study)
 (sertraline metabolite, serotonin uptake by brain inhibition by)
 RN 87857-41-8 CAPLUS
 CN 1-Naphthalenamine, 4-[(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-(9CI) (CA INDEX NAME)

Absolute stereochemistry.




AB Sertraline (I) [79617-96-2] was highly selective and potent competitive inhibitor of synaptosomal serotonin [50-67-9] uptake. I also selectively reduced ex vivo uptake of I and strongly antagonized the serotonin-depleting action of p-chloroamphetamine, indicating potent blockade of serotonin uptake in vivo. Acute and repeated dosing of I decreased serotonin content of whole blood. I only weakly inhibited rat heart uptake of i.v. 3H-labeled norepinephrine [51-41-2]. In substantiation of selective blockade of serotonin uptake, I potentiated various symptoms of 5-hydroxytryptophan but did not reverse reserpine-induced hypothermia. I was a very weak inhibitor of [3H]quinuclidinyl benzilate binding to rat brain membranes in vitro and did not produce anticholinergic effects in mice in vivo. I was well tolerated in mice, rats and dogs, with no locomotor stimulant effects in rats or untoward cardiovascular effects in dogs. The major metabolite, N-desmethylsertraline [87857-41-8] was also a selective serotonin uptake blocker. I strongly reduced immobility of mice in the Porsolt swim test for antidepressants. After repeated dosing in rats, I diminished the cyclic AMP [50-92-4] response of limbic forebrain adenylate cyclase [9012-42-4] to norepinephrine, as well as the binding of [3H]dihydroalprenolol to cortical membranes. It is proposed that selective blockade of serotonin reuptake can induce activation of norepinephrine neurons and subsequent down-regulation of norepinephrine receptors and that I, a highly selective inhibitor of serotonin uptake, may be an efficacious antidepressant without anticholinergic or cardiovascular side-effects.

ACCESSION NUMBER: 1984:17488 CAPLUS
 DOCUMENT NUMBER: 100:17488
 TITLE: Sertraline, 1S,4S-methyl-4-[(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthylamine, a new uptake inhibitor with selectivity for serotonin
 AUTHOR(S): Koe, B. Kenneth; Weissman, Albert; Welch, Willard M.; Browne, Ronald G.
 CORPORATE SOURCE: Cent. Res. Div., Pfizer Inc., Groton, CT, 06340, USA
 SOURCE: Journal of Pharmacology and Experimental Therapeutics (1983), 226(3), 686-700
 CODEN: JPETAB; ISSN: 0022-3565
 DOCUMENT TYPE: Journal
 LANGUAGE: English

L4 ANSWER 79 OF 79 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)
IT 87857-41-8
RL: BIOL (Biological study)
(as sertraline metabolite, serotonin uptake inhibition by,
antidepressant activity in relation to)
RN 87857-41-8 CAPLUS
CN 1-Naphthalenamine, 4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-, (1S,4S)-
(9CI) (CA INDEX NAME)

Absolute stereochemistry.



=> fil reg
COST IN U.S. DOLLARS

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

CA SUBSCRIBER PRICE

SINCE FILE ENTRY	TOTAL SESSION
380.88	536.51

SINCE FILE ENTRY	TOTAL SESSION
-55.30	-55.30

FILE 'REGISTRY' ENTERED AT 21:51:39 ON 14 NOV 2004
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2004 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 12 NOV 2004 HIGHEST RN 780001-49-2
DICTIONARY FILE UPDATES: 12 NOV 2004 HIGHEST RN 780001-49-2



TSCA INFORMATION NOW CURRENT THROUGH MAY 21, 2004

Please note that search-term pricing does apply when conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>
Uploading C:\Program Files\Stnexp\Queries\10663173.str



chain nodes :

17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35

ring nodes :

```

1  2  3  4  5  6  7  8  9 10 11 12 13 14 15 16
chain bonds :
1-25 2-24 3-23 4-22 7-18 8-28 9-27 10-16 11-29 12-17 13-21 13-32 14-19
14-20 15-30 15-31 16-26 32-33 32-34 34-35
ring bonds :
1-2 1-6 2-3 3-4 4-5 5-6 5-13 6-16 7-8 7-12 8-9 9-10 10-11 11-12 13-14
14-15 15-16
exact/norm bonds :
5-13 6-16 13-14 13-32 14-15 15-16 32-34 34-35
exact bonds :
1-25 2-24 3-23 4-22 7-18 8-28 9-27 10-16 11-29 12-17 13-21 14-19 14-20
15-30 15-31 16-26 32-33
normalized bonds :
1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-12 8-9 9-10 10-11 11-12

```

```

Match level :
1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:CLASS 18:CLASS 19:CLASS
20:CLASS 21:CLASS 22:CLASS 23:CLASS 24:CLASS 25:CLASS 26:CLASS 27:CLASS
28:CLASS 29:CLASS 30:CLASS 31:CLASS 32:CLASS 33:CLASS 34:CLASS 35:CLASS

```

L5 STRUCTURE UPLOADED

=> d query

L5 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Structure attributes must be viewed using STN Express query preparation.

```

=> s 15
SAMPLE SEARCH INITIATED 21:53:00 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 677 TO ITERATE

```

```

100.0% PROCESSED        677 ITERATIONS                    0 ANSWERS
SEARCH TIME: 00.00.01

```

```

FULL FILE PROJECTIONS:  ONLINE  **COMPLETE**
                         BATCH  **COMPLETE**
PROJECTED ITERATIONS:            11979 TO        15101
PROJECTED ANSWERS:                0 TO            0

```

L6 0 SEA SSS SAM L5

```

=> s 15 full
FULL SEARCH INITIATED 21:53:04 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 13751 TO ITERATE

```

```

100.0% PROCESSED        13751 ITERATIONS                  5 ANSWERS
SEARCH TIME: 00.00.01

```

L7 5 SEA SSS FUL L5

=> fil caplus

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
155.84	692.35

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
0.00	-55.30

CA SUBSCRIBER PRICE

FILE 'CAPLUS' ENTERED AT 21:53:07 ON 14 NOV 2004

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 14 Nov 2004 VOL 141 ISS 21
FILE LAST UPDATED: 12 Nov 2004 (20041112/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s 17

L8 2 L7

=> d 18 1-2 abs ibib hitstr


L8 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2004 ACS ON STN
 AB Trans-, (1R,4S)-, and
 (1S,4R)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-
 naphthalenamines are claimed. In functional monoamine uptake inhibition
 assays for serotonin, norepinephrine, and dopamine, the (1R,4S)- isomer
 showed IC50 = 0.0077, 0.0096, and 0.0064 µM, resp.
 ACCESSION NUMBER: 2004:252469 CAPLUS
 DOCUMENT NUMBER: 140:287185
 TITLE: Preparation of trans-4-(3,4-dichlorophenyl)-1,2,3,4-
 tetrahydro-1-naphthalenamine for treatment of CNS
 disorders.
 INVENTOR(S): Jerussi, Thomas P.; Fang, Qun Kevin; Currie, Mark
 PATENT ASSIGNEE(S): Sepracor, Inc., USA
 SOURCE: PCT Int. Appl., 36 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004024669	A1	20040325	WO 2003-US29110	20030916
W: RE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RW, GE, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
US 2004092605	A1	20040513	US 2003-663173	20030916
PRIORITY APPLN. INFO.:			US 2002-411304P	P 20020916
			US 2002-411305P	P 20020916

OTHER SOURCE(S): CASREACT 140:287185
 IT 675126-11-1 675126-12-2
 RL: RCT (Reactant); RACT (Reactant or reagent)
 (preparation of dichlorophenyltetrahydronaphthalenamines for
 treatment of
 CNS disorders)
 RN 675126-11-1 CAPLUS
 CN Formamide, N-[(1R,4S)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-
 naphthalenyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

L8 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



RN 675126-12-2 CAPLUS
 CN Formamide, N-[(1S,4S)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-
 naphthalenyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.



IT 674768-11-7P 674768-15-1P 674768-49-1P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT
 (Reactant or reagent)
 (preparation of dichlorophenyltetrahydronaphthalenamines for
 treatment of
 CNS disorders)
 RN 674768-11-7 CAPLUS
 CN Formamide, N-[(1R,4R)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-
 naphthalenyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

L8 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)




RN 674768-15-1 CAPLUS
 CN Formamide, N-[(1S,4R)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-
 naphthalenyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 674768-49-1 CAPLUS
 CN Formamide, N-[(1R,4R)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-
 naphthalenyl]- (9CI) (CA INDEX NAME)

L8 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS
 RECORD. ALL CITATIONS AVAILABLE IN THE RE
 FORMAT

L8 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2004 ACS ON STN
AB Treatment of central nervous system disorders with (1R,4S)-trans- (I) and (1S,4R)-trans-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-N-methyl-1-naphthalenamine (II) is disclosed. I and II were prepared by treating 4-(3,4-dichlorophenyl)tetralinone (III) with (R)-Me3CS(O)NH2 to give the imines which were separated, hydrolyzed to (R)-III and (S)-III, treated

with HCONH2 to give the formamides, which were separated by flash chromatog. and reduced with BH3 to give I and II. I and II had IC50 for 5-HT uptake of 0.0075 and 0.012 μ M, resp.

ACCESSION NUMBER: 2004:252329 CAPLUS

DOCUMENT NUMBER: 140:270634

TITLE: Treatment of CNS disorders with trans-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-N-methyl-1-naphthalenamine

INVENTOR(S): Jerussi, Thomas P.; Fang, Qun Kevin; Currie, Mark

PATENT ASSIGNEE(S): Septacor, Inc., USA

SOURCE: PCT Int. Appl., 24 pp.

DOCUMENT TYPE: CODEN: PIXXD2

LANGUAGE: Patent

FAMILY ACC. NUM. COUNT: English

PATENT INFORMATION: 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004024130	A2	20040325	WO 2003-US29112	20030916
WO 2004024130	A3	20040715		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
US 2004087661	A1	20040506	US 2003-662997	20030916

PRIORITY APPLM. INFO.: US 2002-411303P P 20020916

OTHER SOURCE(S): CASREACT 140:270634; MARPAT 140:270634

IT 674768-11-7P 674768-15-1P 674768-49-1P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)

(preparation of (1R,4S)-trans- and


(1S,4R)-trans-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-N-methyl-1-naphthalenamine for treatment of CNS disorders)

RN 674768-11-7 CAPLUS

CN Formamide, N-[(1R,4R)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenyl]- (9CI) (CA INDEX NAME)


Absolute stereochemistry.

L8 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)




RN 674768-15-1 CAPLUS
CN Formamide, N-[(1S,4R)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenyl]- (9CI) (CA INDEX NAME)

Absolute stereochemistry.



RN 674768-49-1 CAPLUS
CN Formamide, N-[(1R,4R)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenyl]- (9CI) (CA INDEX NAME)

L8 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2004 ACS ON STN (Continued)



=> fil reg
COST IN U.S. DOLLARS

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

CA SUBSCRIBER PRICE

SINCE FILE ENTRY	TOTAL SESSION
10.84	703.19

SINCE FILE ENTRY	TOTAL SESSION
-1.40	-56.70

FILE 'REGISTRY' ENTERED AT 21:55:02 ON 14 NOV 2004
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2004 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 12 NOV 2004 HIGHEST RN 780001-49-2
DICTIONARY FILE UPDATES: 12 NOV 2004 HIGHEST RN 780001-49-2



TSCA INFORMATION NOW CURRENT THROUGH MAY 21, 2004

Please note that search-term pricing does apply when conducting SmartSELECT searches.

Crossover limits have been increased. See HELP CROSSOVER for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=>
Uploading C:\Program Files\Stnexp\Queries\10663173.str



chain nodes :

17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35

ring nodes :

```

1  2  3  4  5  6  7  8  9 10 11 12 13 14 15 16
chain bonds :
1-25 2-24 3-23 4-22 7-18 8-28 9-27 10-16 11-29 12-17 13-21 13-32 14-19
14-20 15-30 15-31 16-26 32-33 32-34 34-35
ring bonds :
1-2 1-6 2-3 3-4 4-5 5-6 5-13 6-16 7-8 7-12 8-9 9-10 10-11 11-12 13-14
14-15 15-16
exact/norm bonds :
5-13 6-16 13-14 13-32 14-15 15-16 32-34 34-35
exact bonds :
1-25 2-24 3-23 4-22 7-18 8-28 9-27 10-16 11-29 12-17 13-21 14-19 14-20
15-30 15-31 16-26 32-33
normalized bonds :
1-2 1-6 2-3 3-4 4-5 5-6 7-8 7-12 8-9 9-10 10-11 11-12

```

```

Match level :
1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:Atom 8:Atom 9:Atom 10:Atom
11:Atom 12:Atom 13:Atom 14:Atom 15:Atom 16:Atom 17:CLASS 18:CLASS 19:CLASS
20:CLASS 21:CLASS 22:CLASS 23:CLASS 24:CLASS 25:CLASS 26:CLASS 27:CLASS
28:CLASS 29:CLASS 30:CLASS 31:CLASS 32:CLASS 33:CLASS 34:CLASS 35:CLASS

```

L9 STRUCTURE UPLOADED

=> d query

L9 STR

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

Structure attributes must be viewed using STN Express query preparation.

=> s l9

SAMPLE SEARCH INITIATED 21:55:31 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 677 TO ITERATE

100.0% PROCESSED 677 ITERATIONS
SEARCH TIME: 00.00.01

0 ANSWERS

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 11979 TO 15101
PROJECTED ANSWERS: 0 TO 0

L10 0 SEA SSS SAM L9

=> s l9 full

FULL SEARCH INITIATED 21:55:34 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 13751 TO ITERATE

100.0% PROCESSED 13751 ITERATIONS
SEARCH TIME: 00.00.01

11 ANSWERS

L11 11 SEA SSS FUL L9

=> fil caplus

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
FULL ESTIMATED COST	ENTRY	SESSION
	155.42	858.61
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
CA SUBSCRIBER PRICE	ENTRY	SESSION
	0.00	-56.70

FILE 'CAPLUS' ENTERED AT 21:55:38 ON 14 NOV 2004
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 14 Nov 2004 VOL 141 ISS 21
 FILE LAST UPDATED: 12 Nov 2004 (20041112/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l11

L12 7 L11

=> d his

(FILE 'HOME' ENTERED AT 21:44:25 ON 14 NOV 2004)

FILE 'REGISTRY' ENTERED AT 21:44:32 ON 14 NOV 2004

L1 STRUCTURE UPLOADED

L2 0 S L1

L3 12 S L1 FULL

FILE 'CAPLUS' ENTERED AT 21:44:55 ON 14 NOV 2004

L4 79 S L3

FILE 'REGISTRY' ENTERED AT 21:51:39 ON 14 NOV 2004

L5 STRUCTURE UPLOADED

L6 0 S L5

L7 5 S L5 FULL

FILE 'CAPLUS' ENTERED AT 21:53:07 ON 14 NOV 2004

L8 2 S L7

FILE 'REGISTRY' ENTERED AT 21:55:02 ON 14 NOV 2004

L9 STRUCTURE UPLOADED

L10 0 S L9

L11 11 S L9 FULL

FILE 'CAPLUS' ENTERED AT 21:55:38 ON 14 NOV 2004


L12 7 S L11

=> s l12 not l8

L13 5 L12 NOT L8

=> d l13 1-5 abs ibib hitstr

L13 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2004 ACS on STN
GI



AB An ortho-substituted BIPHEP ligand, o-Ph-hexaMeO-BIPHEP (I), is designed and synthesized. Compared with chiral biaryl phosphines without ortho substituents such as BINAP and MeO-BIPHEP, I shows higher enantioselectivities in Rh-catalyzed hydrogenation of cyclic enamides.

ACCESSION NUMBER: 2002:308267 CAPLUS
DOCUMENT NUMBER: 137:46985
TITLE: An ortho-Substituted BIPHEP Ligand and Its Applications in Rh-Catalyzed Hydrogenation of Cyclic Enamides

AUTHOR(S): Tang, Wenjun; Chi, Yongxiang; Zhang, Xumu
CORPORATE SOURCE: Department of Chemistry, The Pennsylvania State University, University Park, PA, 16802, USA
SOURCE: Organic Letters (2002), 4(10), 1695-1698
CODEN: ORLEF7; ISSN: 1523-7060

PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English


OTHER SOURCE(S): CASREACT 137:46985
IT 438456-02-1P 438456-04-3P

RL: SPN (Synthetic preparation); PREP (Preparation)
(an ortho-substituted BIPHEP ligand and its applications in Rh-catalyzed hydrogenation of cyclic enamides)

RN 438456-02-1 CAPLUS
CN Carbamic acid, [(1S,4R)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenyl]-, methyl ester (9CI) (CA INDEX NAME)


Absolute stereochemistry. Rotation (-).

L13 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



RN 438456-04-3 CAPLUS
CN Carbamic acid, [(1S,4S)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenyl]-, methyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry.



REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT

L13 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2004 ACS on STN

AB Intramol. endo-cyclization reactions of N-acyliminium ions have seen wide application for the synthesis of heterocyclic compds. The corresponding exocyclic variant, which would provide 1-aminotetralin derivs., for example, has little precedent. The authors have discovered that acyclic N-acylcarbamates can be readily reduced to the corresponding N-acylhemiaminal derivs. in high yield using DIBAL as the reducing agent. These intermediates are remarkably stable and, if desired, can be purified and stored. The acyclic N-acylhemiaminals undergo both intra- and intermol. nucleophilic addition reactions mediated by strong Lewis acids, such as TiCl4. Diastereoselectivity, induced either by a substituent on the newly formed ring, or by utilizing a chiral ester on the carbamic acid, was disappointingly low. This methodol. was successfully applied

to the synthesis of the racemic form of the marketed antidepressant sertraline.

ACCESSION NUMBER: 2001:695659 CAPLUS
DOCUMENT NUMBER: 136:19916
TITLE: The Preparation and Intra- and Intermolecular Addition

Reactions of Acyclic N-Acylimines: Application to the Synthesis of (+)-Sertraline
DeNinno, Michael P.; Eller, Cynthia; Etienne, John B.
PGRD Groton Laboratories, Pfizer Inc., Groton, CT, 06340, USA
Journal of Organic Chemistry (2001), 66(21),

CODEN: JOCEAH; ISSN: 0022-3263
PUBLISHER: American Chemical Society
DOCUMENT TYPE: Journal
LANGUAGE: English


OTHER SOURCE(S): CASREACT 136:19916
IT 374778-34-4P

RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
(preparation and intra- and intermol. addition reactions of acyclic N-acylimines)

RN 374778-34-4 CAPLUS
CN Carbamic acid, [(1R,4R)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenyl]-, methyl ester, rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.

L13 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2004 ACS on STN (Continued)



IT 374778-36-6P
RL: SPN (Synthetic preparation); PREP (Preparation)
(preparation and intra- and intermol. addition reactions of acyclic N-acylimines)
RN 374778-36-6 CAPLUS
CN Carbamic acid, [(1R,4S)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenyl]-, methyl ester, rel- (9CI) (CA INDEX NAME)


Relative stereochemistry.



REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE REFORMAT


L13 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2004 ACS on STN
 AB N-(tert-butoxycarbonyl) amines are directly obtained from oximes of aldehydes and ketones in one-pot when treated with polymethylhydrosiloxane (PMHS) and di-tert-Bu dicarbonate in the presence of catalytic amount of 10% Pd-C for the first time.

ACCESSION NUMBER: 2000:700758 CAPLUS
 DOCUMENT NUMBER: 134:4524
 TITLE: One-pot conversion of oximes to N-(tert-butoxycarbonyl) amines with polymethylhydrosiloxane, Pd-C and di-tert-butyl dicarbonate
 AUTHOR(S): Chandrasekhar, S.; Reddy, M. Venkat; Chandralah, L.
 CORPORATE SOURCE: Indian Institute of Chemical Technology, Hyderabad, 500 007, India
 SOURCE: Synlett (2000), (9), 1351-1353
 CODEN: SYNLES; ISSN: 0936-5214
 PUBLISHER: Georg Thieme Verlag
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 134:4524
 IT 296778-54-6P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (one-pot conversion of oximes to N-(tert-butoxycarbonyl) amines with polymethylhydrosiloxane, Pd-C, and di-tert-Bu dicarbonate)
 RN 296778-54-6 CAPLUS
 CN Carbamic acid,
 [4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenyl]-
 , 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)



REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT


L13 ANSWER 4 OF 5 CAPLUS COPYRIGHT 2004 ACS on STN
 AB One step direct conversion of azides and benzyl carbamates to t-Bu carbamates is achieved using inexpensive and safe hydride source polymethylhydrosiloxane (PMHS) under Pd-C catalysis.
 ACCESSION NUMBER: 2000:488616 CAPLUS
 DOCUMENT NUMBER: 133:266272
 TITLE: Direct conversion of azides and benzyl carbamates to t-butyl carbamates using polymethylhydrosiloxane and Pd-C
 AUTHOR(S): Chandrasekhar, S.; Chandralah, L.; Reddy, Ch. Raji; Reddy, M. Venkat
 CORPORATE SOURCE: Indian Institute of Chemical Technology, Hyderabad, 500 007, India
 SOURCE: Chemistry Letters (2000), (7), 780-781
 CODEN: CMLTAG; ISSN: 0366-7022
 PUBLISHER: Chemical Society of Japan
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 133:266272
 IT 296778-54-6P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (direct conversion of azides and benzyl carbamates to t-Bu carbamates using polymethylhydrosiloxane and Pd-C)
 RN 296778-54-6 CAPLUS
 CN Carbamic acid,
 [4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenyl]-
 , 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)



REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L13 ANSWER 5 OF 5 CAPLUS COPYRIGHT 2004 ACS on STN
 AB An efficient and practical total synthesis of cis-(+)-sertraline is developed involving intramolecular Friedel-Crafts cyclization of an appropriately tailored D-phenylglycine.
 ACCESSION NUMBER: 2000:147120 CAPLUS
 DOCUMENT NUMBER: 132:334264
 TITLE: An expedient total synthesis of cis-(+)-sertraline from D-phenylglycine
 AUTHOR(S): Chandrasekhar, S.; Reddy, M. Venkat
 CORPORATE SOURCE: Indian Institute of Chemical Technology, Hyderabad, 500007, India
 SOURCE: Tetrahedron (2000), 56(8), 1111-1114
 CODEN: TETRAH; ISSN: 0040-4020
 PUBLISHER: Elsevier Science Ltd.
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 OTHER SOURCE(S): CASREACT 132:334264
 IT 267884-84-4P
 RL: RCT (Reactant); SPN (Synthetic preparation); PREP (Preparation); RACT (Reactant or reagent)
 (total synthesis of sertraline from phenylglycine)
 RN 267884-84-4 CAPLUS
 CN Carbamic acid, [(1S,4S)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-1-naphthalenyl]-, 1,1-dimethylethyl ester (9CI) (CA INDEX NAME)

Absolute stereochemistry. Rotation (-).



REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> logoff y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

25.12

883.73

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-3.50

-60.20

STN INTERNATIONAL LOGOFF AT 21:57:41 ON 14 NOV 2004